

## GEOGRAPHICAL DISTRIBUTION.

---

### LIST OF LOCALITIES AT WHICH MACRURA WERE OBTAINED, WITH THE SPECIES TAKEN AT EACH.

STATION VII<sup>P</sup>. February 10, 1873; lat. 28° 35' N., long 16° 5' W.; off Gomera, Canary Islands; depth, 78 fathoms; bottom, volcanic sand. Dredged.

*Arctus pygmæus* (1 ♀).

STATION 13. March 4, 1873; lat. 21° 38' N., long. 44° 39' W.; Mid-north Atlantic; depth, 1900 fathoms; bottom, Globigerina ooze; bottom temperature, 36°·8. Dredged.

*Willemasia leptodactyla* (1 ♀). | *Bentheocaris stylorostratis* (2 ♀ ♂).

STATION 23. March 15, 1873; lat. 18° 24' N., long. 63° 28' W.; off Sombrero Island, West Indies; depth, 450 fathoms; bottom, Pteropod ooze. Dredged.

*Callianassa occidentalis* (1). | *Thaumastocheles zaleuca* (2).

*Cheramus occidentalis* (1). | *Polycheles crucifera* (1).

*Benthesicymus pleocanthus* (1 ♀).

STATION 24. March 25, 1873; lat. 18° 38' 30" N., long. 65° 5' 30" W.; off Culebra Island; depth, 390 fathoms; bottom, Pteropod ooze. Dredged.

*Platybema rugosum* (1 ♀). | *Leptochela serratorbita* (1).

St. Thomas, West Indies; shallow water.

*Sicyonia carinata* (1). | *Alpheus bermudensis* (1 ♀).

*Leptochela serratorbita* (1).

Bermuda; shallow water.

*Stenopus hispidus*. | *Alpheus bermudensis* (3 ♀ ♂).