

Between Bermuda and the Azores. Surface.

*Gennadas intermedius.* | *Brachycarpus savignyii.*

North Atlantic. April 1873. Surface.

<i>Mastigopus suhmi.</i>	<i>Latreutes ensiferus</i> (100). On Gulf-weed.
<i>Sergestes edwardsii.</i>	<i>Hippolyte bidentatus</i> (2 ♂ ♀). On Gulf-weed.
<i>Sergestes atlanticus.</i>	<i>Palæmon natator</i> (113). On Gulf-weed.
<i>Sergestes ovatoculus.</i>	
<i>Sergestes penerinkii.</i>	
<i>Lucifer typus.</i>	
<i>Lucifer reynaudii.</i>	

STATION 40. April 28, 1873; lat. 34° 51' N., long. 68° 30' W.; depth, 2675 fathoms; bottom, blue mud. Dredged.

*Acanthephyra purpurea* (1). | *Acanthephyra sicca* (1).

STATION 42. April 30, 1873; lat. 35° 58' N., long. 70° 35' W.; surface temperature, 65°. Surface.

*Sergestes atlanticus* (1).

STATION 45. May 3, 1873; lat. 38° 34' N., long. 72° 10' W.; south-east of New York; depth, 1240 fathoms; bottom, blue mud; bottom temperature, 37°·2. Dredged.

*Gennadas parvus* (1 ♂). | *Orphania tenuimana* (1).

STATION 49. May 20, 1873; lat. 43° 3' N., long. 63° 39' W.; south of Halifax; depth, 85 fathoms; bottom, gravel, stones; bottom temperature, 35°. Dredged.

<i>Sabinea septemcarinata</i> (22 ♀ ♂).	<i>Hetairus gaimardius</i> (1 ♀).
<i>Hippolyte projecta</i> (1 ♂).	<i>Hetairus tenuis</i> (1 ♂).
<i>Spirontocaris spinus</i> (many ♂ ♀).	<i>Hetairus debilis</i> (15).
<i>Pandalus fulcipes</i> (2 ♀ ♂).	

STATION 57. May 30, 1873; lat. 32° 11' 7" N., long. 65° 3' 20" W.; off Bermuda; depth, 690 fathoms. Dredged.

*Nephropsis rosea.*

STATIONS 62 and 63. June 18, 19, 1873; lat. 35° 7' to 35° 29' N., long. 50° 53' to 52° 32' W. Surface, among Gulf-weed.

<i>Sergestes atlanticus</i> (3 ♂ ♀).	<i>Hippolyte bidentatus</i> (1 ♀). Female with ova.
--------------------------------------	---