

STATION 281. October 6, 1875; lat.  $22^{\circ} 21'$  S., long.  $150^{\circ} 17'$  W.; depth, 2385 fathoms; bottom, red clay; bottom temperature,  $34^{\circ}.9$ . Trawled.

*Haliporus curvirostris* (1)

South Pacific; lat.  $24^{\circ}$  S., long.  $148^{\circ}$  W.

*Sergestes fermerinkii.*

STATION 285. October 14, 1875; lat.  $32^{\circ} 36'$  S., long.  $137^{\circ} 43'$  W.; depth, 2375 fathoms; bottom, red clay; bottom temperature,  $35^{\circ}$ . Trawled.

<i>Haliporus curvirostris</i> (1 ♀).		<i>Bentheocaris exuens</i> (1).
<i>Benthescymus brasiliensis</i> (1 ♀).		<i>Acanthephyra longidens</i> (1 ♂).

STATION 289. October 23, 1875; lat.  $39^{\circ} 41'$  S., long.  $131^{\circ} 23'$  W.; depth, 2550 fathoms; bottom, red clay; bottom temperature,  $34^{\circ}.8$ . Trawled.

<i>Hepomadus inermis</i> (1).		<i>Gennadas parvus</i> (1 ♂).
-------------------------------	--	-------------------------------

South-west Pacific.

*Sergestes junceus.*

STATION 295. November 5, 1875; lat.  $38^{\circ} 7'$  S., long.  $94^{\circ} 4'$  W.; depth, 1500 fathoms; bottom, Globigerina ooze; bottom temperature,  $35^{\circ}.3$ . Trawled.

*Sergestes longicollus* (1).

STATION 298. November 17, 1875; lat.  $34^{\circ} 7'$  S., long.  $73^{\circ} 56'$  W.; depth, 2225 fathoms; bottom, blue mud; bottom temperature,  $35^{\circ}.6$ . Trawled.

*Willemæsia leptodactyla* (♀).

Between Valparaiso and Juan Fernandez. Tow-net at 200 fathoms.

*Sergestes nasidentatus.*

STATION 300. December 17, 1875; lat.  $33^{\circ} 42'$  S., long.  $78^{\circ} 18'$  W.; off Valparaiso; depth, 1375 fathoms; bottom, Globigerina ooze; bottom temperature,  $35^{\circ}.5$ . Trawled.

<i>Pentacheles lavis</i> (1 ♀).		<i>Sergestes profundus</i> (1).
<i>Willemæsia leptodactyla</i> (1 ♂).		<i>Glypocrangon rimapes</i> (2 ♀).
<i>Nematocarcinus proximatus</i> (3 ♀).		