III H	almonio	—continued.	B.G
111. 119			PAGE AND PLATE
in .	Ine	dermal pinuli have moderately long lateral teeth on	
		the long strongly developed radial ray. The large	
		amphidiscs have eight somewhat broad teeth on the	
		margin of the discs,	Hyalonema gracile, . 196
	0.00	22 2 2	XXVII.
	12 The	small parenchymal oxyhexacts have oblique in-	
		wardly directed lateral spines or barbs on their rays,	13
	The	rays of the small parenchymal oxyhexacts are quite	
		smooth,	14
	13 The	body has a cup-like or truncated oval form, with	
		wide oscular aperture, without distinct sieve-	
		membrane. The large amphidiscs have broad,	
		hemispherical arched terminal discs with only six	
		broad terminally lancet-shaped marginal rays,	Hyalonema apertum, . 214
			XXXVII., XXXVIII.
	The	body is spherical or flat and cake-shaped, with narrow	
		oscular aperture, without distinct sieve-membrane.	
		The large amphidiscs have campanulate terminal	
		discs with eight slender pointed rays,	Hyalonema depressum, . 217
		Politica injury	
	14 The	larger, though still comparatively small amphidiscs,	XXXV., XXXVI.
	11 1110	have campanulate terminal discs with eight rather	
		long rays, whose pointed ends reach almost to the	
		middle,	W1.
		initiate,	Hyalonema elegans, . 223
	Th.	James annulistic of the state o	XXXI.
	1116	larger amphidiscs have an approximately spherical	
		form. The hemispherical terminal umbels exhibit	
		eight to twelve broad terminally rounded marginal	
		rays, which often meet, or alternate with those of	77
		the opposite side,	Hyalonema tenerum, 224
			XXXI.
IV.	Nu	merous radially projecting pleuralia (Pheronema,	
		Leidy),	V.
	Wi	thout radial pleuralia. External surface smooth	
2 2027 - 122	<b>2</b> 3.	(Poliopogon, Wyv. Thomson),	VI.
V. P.		a, Leidy.	
		ly markedly longer than broad,	1
		ly not longer than broad,	3
	1 Boo	ly egg- or pear-shaped, narrowed superiorly. The	
		basal tuft consists of several isolated tufts of needles,	Pheronema annæ, .239
			XLII.
	Boo	ly resembles a somewhat broad ellipsoid. The broad	
		basal tuft is bushy,	2
	2 Par	callel to the cuff-like wreath of marginalia, projecting	
		as a fringe round the oscular margin, there is below	
		the oscular margin a connected annular zone of	
		pleuralia, measuring about a finger's breadth, .	Pheronema carpenteri, . 241
			XIIII.