

- Rigid balls, 17.
 Rio de la Plata, *Hyalonema*, 228, 386.
 Roller-stars, 120.
 Root-tuft, 40.
 Rosettes, 30, 48, 128, 175.
Rosettifera (Carter), 9, 138.
Rossella, Carter, 115, 136, 138, 374.
 „ *antarctica*, Carter, 138, 139, 374 (Pl. lv.).
 „ *philippinensis*, 132.
 „ *velata*, Wyv. Thomson, 143, 375.
 Rossellidæ, 129, 374; bathymetrical distribution, 456, 458; geographical distribution, 436; phylogeny, 494.
 Rudiments of rays, 170.
 Sagami Bay, 3; *Aphrocallistes*, 317, 319; *Chonelasma*, 324; *Eurete*, 296, 297, 393; *Farrea*, 277, 278, 391; *Stylocalyx*, 214, 383.
 Santa Cruz, *Pheronema*, 85; *Regadrella*, 239, 387.
 Sarcobacteriellida, 9.
 Sarco-vitreohexactinellida, 9.
 Sars, *Hyalonema longissimum*, 186; *Hyalonema parvum*, 186.
 Schmidt, O., 8; *Aphrocallistes*, 307, 309; *Asconema*, 115; *Cyrtaulon*, 332; *Dactylocalyx*, 346, 347; *Euplectella*, 62; *Farrea*, 268; *Hertwigia*, 109; *Holtenia pourtalesii*, 236; *Hyalonema kentii*, 207; *Hyalonema sieboldii*, 188; *Lanuginella*, 129; *Lefroyella*, 302; *Margaritella*, 351; Mexican Sponges, 15; *Pheronema*, 245; *Pheronema annæ*, 238; phylogeny, 488, 497; *Regadrella*, 84; *Rhabdodictyum*, 107; *Rhabdoplectella*, 108; *Sympagella*, 119.
 Schultze, M., 6; *Euplectella*, 55; *Hyalonemata*, 182; Lophospongiæ, 184.
Scleroplegma, O. Schmidt, 350, 404.
 „ *conicum*, O. Schmidt, 351, 404 (Pl. ci.).
 „ *laterna*, 16.
Sclerothamnus, Marshall, 337, 401.
Sclerothamnus clausii, Marshall, 339 (Pl. xcvi.).
 Scopulæ, 34, 48.
Scopularia (Carter), 289, 391.
Scopulifera (Carter), 9.
 Scotland, *Asconema*, 372; *Pheronema*, 241, 387.
Scyphia, 8.
 Secondary rays, 30.
 Semper, 3; *Euplectella*, 55; *Eurete*, 289; *Hyalonema schultzei* (= *Semperella*), 185, 259.
Semperella, Gray, 259.
 „ *schultzei*, Semper, 261 (Pls. li., lii.).
 Semperellinæ, 259, 389.
 Setubal, *Hyalonema*, 385.
 Seychelle Islands, *Euplectella*, 366.
 Shetland Islands, *Stylocalyx*, 383.
 Sierra Leone, *Malacosaccus*, 93, 369.
 Sieve-plate, 66, 77, 190, 290.
 Silicea (Bowerbank), 183.
 Skeletal spicules, 27.
 Skeleton, 27; evolution of, 502.
 South Atlantic, 440.
 South Indian Ocean, *Balanites*, 372.
 South Pacific, 440; *Bathylorus*, 155, 376; *Dictyocalyx*, 105, 370; *Hyalostylus*, 371; *Stylocalyx*, 224, 385; *Trachycaulus*, 128, 373.
 South Temperate Zone, distribution, 445.
 South Tropical Zone, distribution, 447.
 Sollas, classification, 10; *Dactylocalyx*, 347.
 Spain, *Aphrocallistes*, 395; *Sympagella*, 372.
 Sperms, 24.
 Sphærohexact, 48.
 Sphærohexasters, 31, 48.
 Spicular sheath, 24.
 Spicules, analysis, 28; anchor, 42, 132; appendicular, 132; arrangement, 40, 41; axes of, 29; birotulate, 193; broom-like, 34; clasps of, 78; comital, 68; compass, 80; cruciform, 186; curvature of, 30; extrinsic, 75, 110; floriform-hexradiate, 59; defensive, 71; formation, 499; lamination, 205; orientation, 502; porrecto-multiradiate, 307; position, 40, 41; raphide-like, 81; rationalised, 501; spindle-shaped, 192; stellate, 109; structure, 27; union, 38.
 Spongicolæ, Gray, 182.
Spongia octancyra, Brandt, 182, 184.
 „ *spinicrux*, Brandt, 182, 184.
 Sponge, dredging, 61.
 Stalk, 169, 171, 172.
Stauractinella, Zittel, 14.
 Staurodermidæ, Zittel, 13.
Stauronemata (Sollas), 10, 347.
 Steere's sponge (*Sclerothamnus*), 337.
Stephoscypheus mirabilis (commensal), 98.
 St. Lucia, *Euplectella*, 77.
 St. Thomas, *Aphrocallistes*, 305; *Chonelasma*, 327; *Farrea*, 277, 288; *Hyalonema*, 201; *Myliusia*, 405.
 St. Vincent, *Aulocystis*, 406.
 Stutchbury, *Dactylocalyx*, 342.
Stylocalyx, n. subgen. (see *Hyalonema*), 211, 383.
 „ *apertum*, n. sp., 214, 383 (Pls. xxxvii., xxxviii.).