

PLATE XXVII.

	PAGE
Figs. 1–13. <i>Hyalonema sieboldi</i> , Gray, . . . . .	190
Fig. 1. A dried specimen without the <i>Palythoa</i> -investment of the basal tuft ; x $\frac{1}{3}$ .	
Fig. 2. Terminal sieve-plate, outer side view ; natural size.	
Fig. 3. Oxydiact with central thickening and barbs at both ends ; x 300.	
Figs. 4–6. Small amphidiscs of different sizes ; x 300.	
Fig. 7. Great amphidisc with knobbed axial beam ; x 300.	
Figs. 8, 9. Middle-sized amphidiscs ; x 300.	
Fig. 10. Oxyhexact with bent echinated rays ; x 300.	
Figs. 11, 12. Autodermal pentact pinules ; x 300.	
Fig. 13. Oxyhexact with straight echinated rays ; x 300.	
Figs. 14–23. <i>Hyalonema gracile</i> , F. E. S., . . . . .	196
Fig. 14. Longitudinal section of a somewhat damaged specimen, preserved in alcohol ; natural size.	
Figs. 15–17. Small amphidiscs of different sizes ; x 300.	
Fig. 18. Principal triact with rough ends ; x 300.	
Fig. 19. Greater amphidisc with echinated axial beam ; x 300.	
Fig. 20. Oxyhexact with straight, smooth rays ; x 300.	
Fig. 21. Autodermal pentact pinulus ; x 300.	
Fig. 22. Autogastral pentact pinulus ; x 300.	
Fig. 23. Oxyhexact with smooth bent rays ; x 300.	