

PLATE. LXXIV.

- | | PAGE |
|--|------|
| Figs. 1-6. <i>Farrea sollasii</i> , F. E. S., | 286 |
| Fig. 1. Spicules of a dry specimen in the natural situation of a transverse section of the tube-wall; $\times 100$ | |
| Fig. 2. Uncinatum with numerous spines; $\times 300$. | |
| Fig. 3. Dermal clavula with hemispherical umbel of the head; $\times 450$. | |
| Figs. 4, 5. Gastral clavulæ with anchor teeth of various length; $\times 450$. | |
| Fig. 6. Discohexaster with slender terminal rays; $\times 450$. | |
| Figs. 7-13. <i>Farrea vosmaeri</i> , F. E. S., | 286 |
| Fig. 7. Spicules of a dry specimen, in their natural situation, of a transverse section of the tube-wall; $\times 100$. | |
| Fig. 8. Oxyhexaster with long, strong, principal, and small, divergent, terminal rays; $\times 450$. | |
| Fig. 9. Discohexaster with long, strong, principal, and small, S-shaped, terminal rays, in perianth-like arrangement; $\times 450$. | |
| Fig. 10. Dermal clavula with flatly bent surface of the umbel of the head; $\times 450$. | |
| Fig. 11. Uncinatum with short spines; $\times 300$. | |
| Fig. 12. Gastral clavula with very long spines; $\times 450$. | |
| Fig. 13. Gastral clavula with flatly bent umbel, with somewhat long teeth; $\times 450$. | |