

PLATE II.

Paranebalia longipes (Willemoes Suhm)—*continued*.

- Fig. 1. Adult female, viewed from above ; magnified about thirty-four diameters.
- Fig. 2. Branchial leg of first pair ; magnified fifty diameters.
- Fig. 3. Branchial leg of fifth pair.
- Fig. 4. Branchial leg of last pair.
- Fig. 5. Pleopod of first pair.
- Fig. 6. Pleopod of second pair.
- Fig. 6*a*. Lateral lobe of inner branch of same ; more highly magnified.
- Fig. 6*b*. One of the chitinous hooks from the same lobe ; magnified three hundred and eighty diameters.
- Fig. 7. Pleopod of last pair.
- Fig. 8. First pair of rudimentary caudal limbs.
- Fig. 9. Rudimentary caudal limb of second pair.
- Fig. 10. Last caudal segment, together with right caudal ramus, seen from below.
- Fig. 11. Embryo taken from the incubatory cavity of an adult female specimen, seen from below ; magnified seventy-eight diameters. *R*, rostral plate ; *O*, eyes ; *a*¹, antennulæ ; *a*², antennæ ; *L*, anterior lip ; *M*, mandibles ; *m*¹, maxillæ of first pair ; *m*², maxillæ of second pair ; *brp*, branchial legs ; *pl*, pleopoda ; *ur*, caudal rami.
- Fig. 12. Same embryo, viewed from the right side. *C*, carapace ; remaining letters as in fig. 11.