

GEOGRAPHICAL AND BATHYMETRICAL DISTRIBUTION.

LIST OF LOCALITIES AT WHICH MONAXONID SPONGES WERE OBTAINED, WITH THE SPECIES OBTAINED AT EACH.

STATION 24. March 25, 1873 ; lat. $18^{\circ} 38' 30''$ N., long. $65^{\circ} 5' 30''$ W.; West Indies ; depth, 390 fathoms ; bottom, Pteropod ooze.

Desmacella annexa.

Bermuda, West Indies.

Tedania digitata, var. *bermudensis*.

STATION 49. May 20, 1873 ; lat. $43^{\circ} 3' 0''$ N., long. $63^{\circ} 39' 0''$ W.; south of Halifax, Nova Scotia ; depth, 85 fathoms ; bottom, gravel, stones ; bottom temperature, $35^{\circ} 0$.

Artemisina suberitoides.

Polymastia robusta.

Tentorium semisuberites.

Stylocordyla stipitata.

Quasillina brevis.

STATION 50. May 21, 1873 ; lat. $42^{\circ} 8' 0''$ N., long. $63^{\circ} 39' 0''$ W.; south of Nova Scotia ; depth, 1250 fathoms ; bottom, blue mud ; bottom temperature, $38^{\circ} 0$.

Tentorium semisuberites.

STATION 73. June 30, 1873; lat. $38^{\circ} 30' 0''$ N., long. $31^{\circ} 14' 0''$ W.; depth, 1000 fathoms ; bottom, Pteropod ooze ; bottom temperature, $39^{\circ} 4$.

Rhizochalina fistulosa (?).

Trichostemma sarsii.

STATION 75. July 2, 1873 ; lat. $38^{\circ} 38' 0''$ N., long. $28^{\circ} 28' 30''$ W.; off the Azores ; depth, 450 fathoms ; bottom, volcanic mud.

Reniera implexa.

Gellius angulatus.

Plocamia coriacea, var. *elegans*.

Suberites carnosus.

Suberites elongatus.

Polymastia agglutinans.