TABLE OF GEOGRAPHICAL AND BATHYMETRICAL RANGE—*continued.*

	North Atlantic (Stations I.–110, and 348–354).	South Atlantic (Stations 111–143, and 317–347).	Southern Ocean (Stations 144–160).	Indo-Australian (Stations 161–165, and 181–220).	North Pacific (Stations 221–270).	South Pacific (Stations 165–180, and 271–308).	Patagonian (Stations 304–316).	0–50 Fathoms.	50–200 Fathoms.	200–1000 Fathoms.	1000–3000 Fathoms.
Esperella parishii,				×				×			
,, *gelatinosa*,				×				×			
,, *magellanica*,							×		×		
,, *murrayi*, n. sp.,				×				×			
,, *porosa*, n. sp.,				×				×			
,, *nuda*, n. sp.,		×						×			
,, *fusca*, n. sp.,		×						×			
,, *arenicola*, n. sp.,				×				×			
,, *simonis*, n. sp.,		×						×			
,, *biserialis*, n. sp.,						×					×
,, sp.,				×				×			
Esperiopsis symmetrica, n. sp.,			×							×	
,, *edwardii*, var. *americana*, nov.,							×		×		
,, *cylindrica*, n. sp.,				×				×			
,, *challengeri*, n. sp.,				×						×	
,, ,, var. *meangensis*, nov.,				×						×	
,, *profunda*, n. sp.,			×								×
,, *anomala*, n. sp.,					×			×			
,, *pulchella*, n. sp.,				×					×		
Cladorhiza abyssicola, var. *rectangularis*, nov.,						×					×
,, ,, var. *linearis*, nov.,						×					×
,, *moruliformis*, n. sp.,			×								×
,, *longipinna*, n. sp.,					×						×
,, *similis*, n. sp.,						×					×
,, *inversa*, n. sp.,		×									×
,, (?) *tridentata*, n. sp.,			×								×
Axoniderma mirabile, n. gen. et sp.,						×					×
Chondrocladia concrescens (?),					×						×
,, *clavata*, n. sp.,						×			×		
,, *crinita*, n. sp.,				×							×
Melliderma stipitata, n. gen. et sp.,						×					×
Desmacidon fruticosa, var.,				×					×		
,, *reptans*, n. sp.,		×						×			
,, *conulosa*, n. sp.,		×						×			
(?) *ramosa*, n. sp.,		×	×						×		