TABLE OF GEOGRAPHICAL AND BATHYMETRICAL RANGE—*continued.*

	North Atlantic (Stations I.–110, and 348–364).	South Atlantic (Stations 111–143, and 317–347).	Southern Ocean (Stations 144–160).	Indo-Australian (Stations 161–165, and 181–220).	North Pacific (Stations 221–270).	South Pacific (Stations 166–180, and 271–303).	Patagonian (Stations 304–316).	0–50 Fathoms.	50–200 Fathoms.	200–1000 Fathoms.	1000–3000 Fathoms.
Axinella (?) *tubulosa*, n. sp.,		×								×	
,, (?) *paradoxa*, n. sp.,		×							×		
Raspailia tenuis, n. sp.,		×						×			
,, *flagelliformis*, n. sp.,		×						×			
,, (?) *rigida*, n. sp.,		×							×		
Dendropsis bidentifera, n. gen. et sp.,		×						×			
Thrinacophora cervicornis, n. gen. et sp.,				×				×			
,, *funiformis*, n. sp.,		×						×			
Suberites carnosus,	×	×		×				×		×	
,, *caminatus*, n. sp.,			×						×		
,, *microstomus*, n. sp.,			×						×		
,, *perfectus*, n. sp.,				×				×			
,, *antarcticus*,			×						×		
,, *axiatus*, n. sp.,		×								×	
,, *durissimus*, n. sp.,				×							
,, *mollis*, n. sp.,			×							×	
,, *elongatus*, n. sp.,	×									×	
,, *spiralis*, n. sp.,							×				
,, *ramulosus*, n. sp.,				×						×	
,, ,, var. *cylindrifera*, nov.,				×					×		
,, *senilis*, n. sp.,					×						×
Polymastia robusta,	×								×		
,, *corticata*, n.		×									×
,, *agglutinans*, n. sp.,	×									×	
Proteleia sollasi, n. gen. et sp.,		×						×			
Trichostemma sarsii, n. sp.,	×			×						×	×
,, *irregulare*, n. sp.,						×					×
Tentorium semisuberites,	×	×							×		×
Stylocordyla stipitata,	×	×	×					×	×		×
,, ,, var. *globosa*, nov.,			×					10–	–100		
Quasillina brevis,	×								×		
Cliona dissimilis, n. sp.,				×				×			
Spirastrella decumbens, var.,				×				×			
,, *massa*, n. sp.,				×				×			
,, *solida*, n. sp.,				×				×			