

PLATE XVII.

	PAGE
Fig. 1. <i>Iophon chelifer</i> ; <i>a, b</i> , spined styli; <i>c</i> , tylote spicule; × 180. . . . .	119
Fig. 2. <i>Vomerula esperioides</i> ; <i>a, b</i> , two views of the diancistron, × 180; <i>c</i> , sigma, × 475; <i>d</i> , young diancistron, showing the hooks still united to the shaft by a thin web, × 475, . . . . .	61
Fig. 3. <i>Iophon chelifer</i> ; <i>a, b, c, d</i> , different views of the bipocilli; × 900, . . . . .	119
Fig. 4. <i>Vomerula esperioides</i> , stylus; × 180, . . . . .	61
Fig. 5. <i>Iophon abnormalis</i> ; <i>a, b</i> , tylota; <i>c</i> , stylus; × 180, . . . . .	122
Fig. 6. <i>Iophon cylindricus</i> ; <i>a, b</i> , styli; <i>c</i> , tylote spicule; × 180, . . . . .	120
Fig. 7. <i>Iophon abnormalis</i> , rosette of anisochelæ; × 540, . . . . .	122
Fig. 8. <i>Iophon chelifer</i> , anisochela, side view; × 1200, . . . . .	119
Fig. 9. <i>Iophon laminalis</i> ; <i>a, b, c</i> , three views of the bipocilli; × 1350, . . . . .	121
Fig. 10. <i>Iophon (?) omnivorus</i> , portion of dermal layer, showing dermal skeleton of foreign bodies, and pores; × 175, . . . . .	122
Fig. 11. <i>Iophon laminalis</i> ; <i>a</i> , tylote spicule; <i>b</i> , stylus; × 180, . . . . .	121
Fig. 12. <i>Vomerula esperioides</i> , portion of dermal layer, showing pores and dermal skeleton; × 63, . . . . .	60