

<i>Actinometra duplex.</i>	<i>Actinometra paucicirra.</i>
<i>echinoptera.</i>	<i>pectinata.</i>
<i>elongata.</i>	<i>peroni.</i>
<i>fimbriata.</i>	<i>quadrata.</i>
<i>grandicalyx.</i>	<i>regalis.</i>
<i>japonica.</i>	<i>robustipinna.</i>
<i>lineata, II.</i>	<i>rotalaria.</i>
<i>littoralis.</i>	<i>rubiginosa.</i>
<i>maculata.</i>	<i>schlegeli.</i>
<i>magnifica.</i>	<i>sentosa.</i>
<i>meridionalis, II.-V.</i>	<i>simplex.</i>
<i>multibrachiata.</i>	<i>solaris.</i>
<i>multifida.</i>	<i>stelligera, V.</i>
<i>multiradiata.</i>	<i>trichoptera.</i>
<i>nigra.</i>	<i>typica, V.</i>
<i>nobilis.</i>	<i>variabilis.</i>
<i>novæ-guineæ.</i>	<i>valida.</i>
<i>parvicirra, V.</i>	

Totals.—*Antedon*, 52 species ; *Actinometra*, 45 species.

TABLE II.—Species found at depths between 20 and 50 fathoms.

<i>Antedon armata.</i>	<i>Antedon prolixa, III.-VII.</i>
<i>carinata, I.-V.</i>	<i>quadrata, III.-VI.</i>
<i>denticulata.</i>	<i>rosacea, I.-III.</i>
<i>elegans, I.</i>	<i>variipinna, I.</i>
<i>eschrichti, I.-VII.</i>	<i>Actinometra lineata, I.</i>
<i>macronema, I.</i>	<i>meridionalis, I.-V.</i>
<i>magellanica.</i>	<i>Eudiocrinus indivisus.</i>
<i>petasus, I.-III.</i>	<i>Promachocrinus kerguelensis, III., IV.</i>
<i>phalangium, III.-V.</i>	

Totals.—*Antedon*, 13 species ; *Actinometra*, 2 species.

TABLE III.—Species found at depths between 50 and 100 fathoms.

<i>Antedon antarctica.</i>	<i>Antedon duplex, IV., V.</i>
<i>balanoides.</i>	<i>eschrichti, I.-VII.</i>
<i>carinata, I.-V.</i>	<i>hageni, IV., V.</i>
<i>compressa, IV.</i>	<i>petasus, I., II.</i>
<i>defecta, IV., V.</i>	<i>phalangium, II.-V.</i>