

TABLE VIII.—Species found at depths between 800 and 1100 fathoms.

<i>Antedon alternata</i> , VII.		<i>Eudiocrinus varians</i> .
<i>spinicirra</i> .		<i>semperi</i> , VII.
Pentacrinoid larvæ of <i>Antedon</i> , sp.		

TABLE IX.—Species found at depths between 1100 and 1500 fathoms.

<i>Antedon acutiradia</i> .		<i>Antedon breviradia</i> , VII.
<i>basicurva</i> <sup>1</sup> ?		<i>inæqualis</i> ?

TABLE X.—Species found at depths between 1500 and 1800 fathoms.

<i>Antedon abyssorum</i> .		<i>Promachocrinus abyssorum</i> .
<i>bispinosa</i> .		<i>Thaumatocrinus renovatus</i> .
<i>remota</i> .		

TABLE XI.—Species found at 2600 and 2900 fathoms.

*Antedon abyssicola*.

## SUMMARY.

## A.

I.	97	species	found	at	depths	down	to	20	fathoms.
II.	17	species	found	between	20	and	50	fathoms.	
III.	20	„	„	50	and	100	„		
IV.	34	„	„	100	and	200	„		
V.	22	„	„	200	and	350	„		
VI.	22	„	„	350	and	500	„		
VII.	19	„	„	500	and	800	„		
VIII.	5	„	„	800	and	1100	„		
IX.	2	„	„	1100	and	1500	„		
X.	5	„	„	1500	and	1800	„		
XI.	1	„	„	2600	and	2900	„		

## B.

- I. 97 species found at depths down to 20 fathoms.  
 I.-II. 9 of these descend to 50 fathoms.  
 I.-III. 5 of which reach 100 fathoms.  
 I., V. 3 also occur at 210 or 255 fathoms.

<sup>1</sup> Only if obtained from 1350 fathoms at Station 175.