

*Stenella*, Gray.

The species of this genus appear to be widely scattered. *Stenella imbricata* (Y. Johns.), the first species described, was taken off Madeira; *Stenella johnsoni* was dredged off Ascension; *Stenella acanthina* occurs off the Rio de la Plata; *Stenella spinosa* was found off Prince Edward Island; *Stenella gigantea* was found at Kandavu, in the Pacific Ocean, while *Stenella doederleini* was dredged off Yokohama, Japan.

*Thouarella*, Gray.

The distribution of the species of this genus in the Atlantic Ocean is limited to the Falkland Islands, where the type specimen of *Thouarella antarctica* (Val.) was found by du Petit Thouars and Capt. J. Clark; to the coast of Patagonia, where *Thouarella köllikeri* was dredged; and to Tristan da Cunha, where *Thouarella affinis* was found. *Thouarella antarctica* (Val.), *Thouarella variabilis* and vars., *Thouarella affinis*, var., have been taken at Prince Edward Island and Heard Island in the Antarctic Ocean. In the Pacific Ocean two species occur, *Thouarella moseleyi* off the Kermadecs, and *Thouarella hilgendorfi* (Std.), which has been met with both south of Papua and at Japan.

*Amphilaphis*, Wright and Studer.

The only species of this genus (*Amphilaphis regularis*) was taken at Tristan da Cunha.

*Plumarella*, Gray.

Of the three species described, one, *Plumarella pourtalesi*, Ver., has been found off the coast of Florida, a second, *Plumarella delicatissima*, off Port Grappler, Patagonia, and the third, *Plumarella penna* (Lamk.), is referred by Milne-Edwards to Australia.

*Caligorgia*, Gray.

The distribution of the species of this genus is rather peculiar; in the Atlantic Ocean *Caligorgia verticillata* (Pall.) is found off the north-west coast of Africa, and extending into the Mediterranean, while *Caligorgia gracilis* (M.-Edw.) is found in the West Indies. In the Indian Ocean, on the authority of a specimen in the Bern Museum, *Caligorgia flabellum* (Ehrb.) occurs at the Mauritius. In the Pacific Ocean *Caligorgia ventilabrum*, Std., is found off the north of New Zealand, *Caligorgia sertosa* south of Papua, *Caligorgia flabellum* (Ehrb.) occurs at Japan and at Formosa, while *Caligorgia compressa* (Ver.) occurs at the Aleutian Islands.