

found in the most recent Tertiary deposits, and have received other titles in that connection.

<i>Creseis acicula</i> , Rang.	<i>Cleodora lessonii</i> , Rang.
<i>Creseis acus</i> , Eschscholtz.	<i>Cleodora lobata</i> , Sowerby.
<i>Cleodora andrææ</i> , Boas.	<i>Cleodora martensii</i> , Pfeffer.
<i>Hyalæa australis</i> , d'Orbigny.	<i>Creseis monotis</i> , Troschel.
<i>Cleodora balantium</i> , Rang.	<i>Cleodora munda</i> , Gould.
<i>Balantium bicarinatum</i> , Benson.	<i>Cleodora obtusa</i> , Quoy and Gaimard.
<i>Cleodora brownii</i> , de Blainville.	<i>Cleodora occidentalis</i> , Dall.
<i>Creseis caligula</i> , Eschscholtz.	<i>Clio pellucidum</i> , Gray.
<i>Clio caudata</i> , Linné.	<i>Creseis phæostoma</i> , Troschel.
<i>Cleodora chaptalii</i> , Souleyet.	<i>Cleodora placida</i> , Gould.
<i>Cleodora chierchiæ</i> , Boas.	<i>Balantium politum</i> , Craven, MS.
<i>Creseis clava</i> , Rang.	<i>Cleodora pygmæa</i> , Boas.
<i>Creseis compressa</i> , Eschscholtz.	<i>Clio pyramidata</i> , Linné.
<i>Cleodora compressa</i> , Souleyet.	<i>Cleodora quadrispinosa</i> , Rang.
<i>Creseis conica</i> , Eschscholtz.	<i>Styliola recta</i> , Lesueur.
<i>Creseis conoidea</i> , A. Costa.	<i>Balantium recurvum</i> , Benson.
<i>Hyalæa corniformis</i> , d'Orbigny.	<i>Clio rugosum</i> , Gray.
<i>Creseis cornucopiæ</i> , Eschscholtz.	<i>Creseis rugulosa</i> , Cantraine.
<i>Cleodora curvata</i> , Souleyet.	<i>Creseis spinifera</i> , Rang.
<i>Hyalæa cuspidata</i> , Bosc.	<i>Creseis striata</i> , Rang.
<i>Clio depressa</i> , Gray.	<i>Creseis striata</i> , Delle Chiaje.
<i>Cleodora exacuta</i> , Gould.	<i>Cleodora subula</i> , Quoy and Gaimard.
<i>Cleodora falcata</i> , Gould.	<i>Cleodora sulcata</i> , Pfeffer.
<i>Cleodora falcata</i> , Pfeffer.	<i>Hyalæa tricuspидata</i> , Bowdich.
<i>Cleodora flexa</i> , Pfeffer.	<i>Cleodora trifilis</i> , Troschel.
<i>Cleodora inflata</i> , Souleyet.	<i>Creseis unguis</i> , Eschscholtz.
<i>Cleodora lamartinieri</i> , Rang.	<i>Creseis virgula</i> , Rang.
<i>Hyalæa lanceolata</i> , Lesueur.	<i>Styliola vitrea</i> , Verrill.
	<i>Creseis zonata</i> , Delle Chiaje.

These fifty-seven names may be classified as follows :—

1. One may be discarded as not referring to a Pteropod, viz., *Creseis rugulosa*, Cantraine,¹ which is really a Gastropod of the genus *Cæcum*.

2. One refers to the adult stage of another genus of Thecosomatous Pteropods, viz., *Cleodora obtusa*, Quoy and Gaimard, which is probably the same as *Cuvierina columnella*.

¹ Malacologie Méditerranéenne et littorale, *Mém. Acad. d. Sci. Bruxelles*, t. xiii. p. 32.