

3. Seven names refer to young stages of the Thecosomata of the genus *Cavolinia*, viz.:—

*Cleodora compressa*, Souleyet = *Cavolinia trispinosa*.

*Cleodora curvata*, Souleyet = *Cavolinia uncinata*.

*Clio depressa*, Gray = *Cleodora compressa*, Souleyet = *Cavolinia trispinosa*.

*Clio pellucidum*, Gray = *Pleuropus pellucidus*, Eschscholtz = *Cavolinia inflexa*.

*Balantium rugosum*, Gray = *Hyalæa rugosa*, d'Orbigny = *Cavolinia gibbosa*.

*Cleodora pygmæa*, Boas = *Cavolinia quadridentata*.

*Cleodora trifilis*, Troschel = *Cavolinia*, sp.

The attentive study of the forms designated by the remaining specific titles shows that thirty-three of these ought to be discarded, and considered only as synonyms of the fourteen remaining species.

<i>Creseis acus</i> , Eschscholtz,	.	= <i>Creseis acicula</i> , Rang.
<i>Balantium bicarinatum</i> , Benson,		= <i>Cleodora balantium</i> , Rang.
<i>Cleodora brownii</i> , de Blainville,		= <i>Clio pyramidata</i> , Linné.
<i>Creseis caligula</i> , Eschscholtz,	.	= <i>Creseis virgula</i> , Rang.
<i>Clio caudata</i> , Linné,	.	= ? <i>Clio pyramidata</i> , Linné.
<i>Creseis clava</i> , Rang,	.	= <i>Creseis acicula</i> , Rang.
<i>Creseis compressa</i> , Eschscholtz,	.	= <i>Creseis striata</i> , Rang.
<i>Creseis conoidea</i> , O. Costa,	.	= <i>Creseis conica</i> , Eschscholtz.
<i>Hyalæa corniformis</i> , d'Orbigny,		= <i>Creseis virgula</i> , Rang.
<i>Creseis cornucopiae</i> , Eschscholtz,		= <i>Creseis virgula</i> , Rang.
<i>Cleodora exacuta</i> , Gould,	.	= <i>Clio pyramidata</i> , Linné.
<i>Cleodora falcata</i> , Gould,	.	= <i>Creseis virgula</i> , Rang.
<i>Cleodora falcata</i> , Pfeffer,	.	= <i>Balantium politum</i> , Craven, MS.
<i>Cleodora flexa</i> , Pfeffer,	.	= <i>Creseis virgula</i> , Rang.
<i>Cleodora inflata</i> , Souleyet,	.	= <i>Cleodora balantium</i> , Rang (young).
<i>Cleodora lamartinieri</i> , Rang,	.	= <i>Clio pyramidata</i> , Linné.
<i>Hyalæa lanceolata</i> , Lesueur,	.	= <i>Clio pyramidata</i> , Linné.
<i>Cleodora lessoni</i> , Rang,	.	= <i>Hyalæa cuspidata</i> , Bosc.
<i>Cleodora lobata</i> , Sowerby,	.	= <i>Clio pyramidata</i> , Linné.
<i>Cleodora martensii</i> , Pfeffer,	.	= <i>Clio pyramidata</i> , Linné.
<i>Creseis monotis</i> , Troschel,	.	= <i>Creseis striata</i> , Rang.
<i>Cleodora munda</i> , Gould,	.	= <i>Creseis virgula</i> , Rang.
<i>Creseis phæostoma</i> , Troschel,	.	= <i>Creseis striata</i> , Rang.
<i>Creseis placida</i> , Gould,	.	= <i>Creseis virgula</i> , Rang.
<i>Cleodora quadrispinosa</i> , Rang,		= <i>Hyalæa cuspidata</i> , Bosc.
<i>Styliola recta</i> , Lesueur,	.	= <i>Cleodora subula</i> , Quoy and Gaimard.