

2. Subgenus *Hyalocylrix*.*Creseis striata*, Rang.3. Subgenus *Styliola*.*Cleodora subula*, Quoy and Gaimard.4. Subgenus *Clio*, s. str.*Balantium politum*, Craven, MS.*Cleodora andreae*, Boas.*Cleodora balantium*, Rang.*Hylæa australis*, d'Orbigny.*Cleodora sulcata*, Pfeffer.*Clio pyramidata*, Linné.*Hyalæa cuspidata*, Bosc.

The four sections may be distinguished as follows :—

I. Shell without lateral keels.

1. Shell without dorsal longitudinal groove.

- | | |
|---|-------------------------|
| A. Shell with a circular section, | 1. <i>Creseis</i> . |
| B. Shell flattened dorso-ventrally, with transverse grooves over its entire length, | 2. <i>Hyalocylrix</i> . |

2. Shell with a dorsal longitudinal groove,

- | | |
|---|---|
| I. Shell without lateral keels, | 3. <i>Styliola</i> . |
| II. Shell with lateral keels, | 4. <i>Clio</i> , including <i>Balantium</i> . |

Subgenus *Creseis*,¹ Rang (s. str.).

1828. *Creseis*, Rang, Notice sur quelques Mollusques nouveaux appartenant au genre *Cleodora*, &c., Ann. d. Sci. Nat., sér. 1, t. xiii. p. 302.

Styliola, Auctorum, non Lesueur.

Characters and Description.—Shell elongated, of conical form, with a circular transverse section, with a smooth surface on its initial portion at least, with the embryonic portion not marked off by a deep constriction, with a rounded apex.

Animal with the left tentacle very rudimentary, the fin exhibiting a small narrow projecting lobe on the proximal half of the dorsal (anterior) margin, the opening of the mantle as broad as that of the shell.

¹ Mythological name.