

(d'Orbigny placed in the genus *Hyalæa* = *Cavolinia* all the members of the family Cavoliniidæ), or even to other groups, and confine our attention to the forms really belonging to this genus, we shall find, for extant species, fifty-four specific titles, of which three are used with several connotations, which increases the mischief still further.

Hyalæa affinis, d'Orbigny.
Hyalæa angulata, Souleyet.
Hyalæa australis, Péron.
Hyalæa chemnitziana, Péron and Lesueur.
Hyalæa complanata, Gegenbaur.
Cleodora compressa, Souleyet.
Hyalæa cornea, Lamarck.
Hyalæa costata, Pfeffer.
Hyalæa cumingii, Sowerby.
Cleodora curvata, Souleyet.
Hyalæa cuspidata, Delle Chiaje (non Bosc).
Hyalæa depressa, Bivona (non d'Orbigny).
Hyalæa depressa, d'Orbigny.
Hyalæa ecaudata, Lesueur.
Hyalæa elongata, Lesueur.
Hyalæa femorata, Gould.
Hyalæa fissirostris, Benson.
Hyalæa flava, d'Orbigny.
Hyalæa forskahlii, Lesueur.
Hyalæa gegenbauri, Pfeffer.
Hyalæa gibbosa, Rang.
Hyalæa globulosa, Rang.
Pleuropus hargeri, Verrill.
Hyalæa imitans, Pfeffer.
Hyalæa inermis, Gould.
Hyalæa inflexa, Lesueur.
Hyalæa intermedia, Sowerby.
Hyalæa labiata, d'Orbigny.
Hyalæa lævigata, d'Orbigny.

Hyalæa limbata, d'Orbigny.
Pleuropus longifilis, Troschel.
Hyalæa longirostris, Lesueur.
Hyalæa minuta, Sowerby.
Hyalæa mucronata, Quoy and Gaimard.
Cavolina natans, Abildgaard.
Hyalæa obtusa, Sowerby.
Hyalæa papilionacea, Quoy and Gaimard.
Pleuropus pellucidus, Eschscholtz.
Hyalæa peroni, Lesueur.
Cavolina pisum, Mørch.
Cleodora pygmæa, Boas.
Hyalæa quadridentata, Lesueur.
Hyalæa quadrispinosa, d'Orbigny.
Hyalæa reeviana, Dunker.
Hyalæa rotundata, Boas.
Hyalæa rugosa, d'Orbigny.
Hyalæa teniobranchea, Péron and Lesueur.
Hyalæa triacantha, Bronn.
Anomia tridentata, Forskål.
Cleodora trifilis, Troschel.
Hyalæa trispinosa, Lesueur.
Hyalæa truncata, Krauss (non Lesueur).
Hyalæa truncata, Lesueur.
Hyalæa uncinata, Hoeninghaus (non Rang).
Hyalæa uncinata, Rang.
Hyalæa uncinatiformis, Pfeffer.

Hyalæa vaginellina, Cantraine.