

GEOGRAPHICAL DISTRIBUTION.

A. STATION LIST OF THE THECOSOMATOUS PTEROPODA OF THE CHALLENGER EXPEDITION.

I. STATIONS WHERE LIVING SPECIMENS WERE COLLECTED BY TRAWLING OR DREDGING.

STATION VIIF. February 2, 1873; off Madeira; lat. 32° 27' 0" N., long. 16° 40' 30" W.; surface temperature, 63° F.

Clio (Creseis) acicula. | *Cavolinia trispinosa.*
Cavolinia globulosa.

On April 14, 1873; off Bermuda; lat. 32° 18' N., long. 64° 53' W.

Cavolinia longirostris.

STATION 53. May 26, 1873; Halifax to Bermuda; lat. 36° 30' N., long. 63° 40' W.; surface temperature, 73° F.

Cuvierina columnella.

STATION 63. June 19, 1873; Bermuda to Azores; lat. 35° 29' N., long. 50° 53' W.; surface temperature, 71° F.

Clio (Creseis) acicula. | *Cuvierina columnella.*
Clio pyramidata. | *Cavolinia trispinosa.*

STATION 81. July 13, 1873; Azores to Madeira; lat. 34° 11' N., long. 19° 52' W.; surface temperature, 71° F.

Clio (Creseis) acicula.

STATION 100. August 16, 1873; St. Vincent to St. Paul's Rocks; lat. 7° 1' N., long. 15° 55' W.; surface temperature, 79° F.

Cavolinia uncinata.