

STATION 159. March 10, 1874 ; Termination land to Melbourne ; lat. 47° 25' S., long. 130° 22' E.; surface temperature, 51°·5 F.

Clio australis.

On March 15, 1874 ; off Melbourne ; lat. 39° 45' S., long. 40° 40' E.; surface temperature, 59° F.

Clio (Styliola) subula.

Clio pyramidata.

On March 16, 1874 ; off Melbourne ; lat. 39° 22' S., long. 142° 22' E.; surface temperature, 62° F.

Clio (Styliola) subula.

Between Stations 162 and 163. April 3, 1874 ; Melbourne to Sydney ; lat. 38° 7' S., long. 149° 18' E.; surface temperature, 65° F.

Limacina inflata.

Clio (Creseis) virgula.

Limacina lesueurii.

Cavolinia trispinosa (young, as
" *Cleodora compressa* ").

Limacina trochiformis.

Cavolinia longirostris.

Limacina bulimoides.

STATION 163. April 4, 1874 ; Melbourne to Sydney ; lat. 36° 57' S., long. 150° 34' E.; surface temperature, 72° F.

Limacina bulimoides.

STATION 164A. June 13, 1874 ; off Sydney ; lat. 34° 9' S., long. 151° 55' E.; surface temperature, 70°·2 F.

Clio (Creseis) virgula.

Cavolinia inflexa (young, as
" *Hyalæa depressa* ").

Clio (Creseis) acicula.

Clio (Styliola) subula.

STATION 175. August 12, 1874 ; Fiji to Raine Island ; lat. 19° 2' S., long. 177° 10' E.; surface temperature, 77°·5 F.

Limacina inflata.

Clio (Styliola) subula.

Limacina lesueurii.

Clio pyramidata.

Limacina bulimoides.

Cuvierina columnella.

Clio (Creseis) acicula.

Cavolinia quadridentata.

Clio (Hyalocylis) striata.

Cavolinia longirostris.

Cavolinia inflexa.