

STATION 216A. February 16, 1875; north of New Guinea; lat. $2^{\circ} 56' N.$, long. $134^{\circ} 11' E.$; surface temperature, $82^{\circ} F.$

| | |
|--------------------------------|---|
| <i>Limacina inflata.</i> | <i>Cuvierina columnella.</i> |
| <i>Limacina lesueuri.</i> | <i>Cavolinia quadridentata.</i> |
| <i>Limacina trochiformis.</i> | <i>Cavolinia longirostris</i> (young, as <i>Hyalæa lavigata</i>). |
| <i>Clio (Creseis) virgula.</i> | <i>Cavolinia globulosa.</i> |
| <i>Clio (Creseis) acicula.</i> | Fry of <i>Cymbulia</i> and <i>Gleba</i> . |
| <i>Clio balantium.</i> | |

Between Stations 229 and 230. April 3, 1875; Admiralty Islands to Yokohama; lat. $24^{\circ} 49' N.$, long. $138^{\circ} 34' E.$; surface temperature, $71^{\circ} F.$

| | |
|--------------------------------|--------------------------------|
| <i>Clio (Creseis) virgula.</i> | <i>Clio (Styliola) subula.</i> |
| | <i>Cavolinia gibbosa.</i> |

Near Station 230. April 4, 1875; Admiralty Islands to Yokohama; lat. $25^{\circ} 33' N.$, long. $137^{\circ} 57' E.$; surface temperature, $69^{\circ} F.$

| | |
|--------------------------------|---------------------------|
| <i>Clio (Styliola) subula.</i> | <i>Cavolinia gibbosa.</i> |
| <i>Clio cuspidata.</i> | <i>Cavolinia inflexa.</i> |

STATION 230. April 5, 1875; Admiralty Islands to Yokohama; lat. $26^{\circ} 29' N.$, long. $137^{\circ} 57' E.$; surface temperature, $69^{\circ} F.$

| | |
|-----------------------------------|--------------------------------|
| <i>Clio (Creseis) virgula.</i> | <i>Cavolinia trispinosa.</i> |
| <i>Clio (Hyalocylix) striata.</i> | <i>Cavolinia longirostris.</i> |
| <i>Clio pyramidata.</i> | <i>Cavolinia gibbosa.</i> |
| <i>Cuvierina columnella.</i> | <i>Cavolinia inflexa.</i> |

On May 4, 1875; at Yokohama; surface temperature, $60^{\circ} F.$

Cavolinia inflexa (young, as "*Pleuropus pellucidus*"; *fide* Willemoes Suhm).

STATION 237. June 17, 1875; off Japan; lat. $34^{\circ} 37' N.$, long. $140^{\circ} 32' E.$; surface temperature, $73^{\circ} F.$

Clio (Creseis) acicula.

STATION 241. June 23, 1875; off Japan; lat. $35^{\circ} 41' N.$, long. $157^{\circ} 42' E.$; surface temperature, $69^{\circ} 2 F.$

Cavolinia tridentata.

STATION 243. June 26, 1875; Yokohama to Sandwich Islands; lat. $35^{\circ} 24' N.$, long. $166^{\circ} 35' E.$; surface temperature, $71^{\circ} F.$

Limacina bulimoides.