

On June 29, 1875 ; Yokohama to Sandwich Islands ; lat. $35^{\circ} 49'$ N., long. $171^{\circ} 46'$ E.; surface temperature, 69° F.

Gleba spectabilis.

Between Stations 247 and 248. July 4, 1875 ; Yokohama to Sandwich Islands ; lat. $36^{\circ} 42'$ N., long. $179^{\circ} 50'$ W.; surface temperature, 70° F.

Limacina lesueuri.

Cavolinia trispinosa (young, as
“*Cleodora compressa*”).

Limacina bulimoides.

Clio (Hyaloclylix) striata.

Cavolinia inflexa (young, as “*Hyalaea depressa*”).

STATION 249. July 7, 1875 ; Yokohama to Sandwich Islands ; lat. $37^{\circ} 59'$ N., long. $171^{\circ} 48'$ W.; surface temperature, $65^{\circ} \cdot 2$ F.

Cavolinia gibbosa.

STATION 251. July 10, 1875 ; Yokohama to Sandwich Islands ; lat. $37^{\circ} 37'$ N., long. $163^{\circ} 26'$ W.; surface temperature, 65° F.

Clio pyramidata.

STATION 254. July 17, 1875 ; Yokohama to Sandwich Islands ; lat. $35^{\circ} 13'$ N., long. $154^{\circ} 43'$ W.; surface temperature, 72° F.

Clio (Hyaloclylix) striata.

Clio cuspidata.

Clio pyramidata.

Cavolinia tridentata.

Cymbulia sp.

STATION 256. July 21, 1875 ; Yokohama to Sandwich Islands ; lat. $30^{\circ} 22'$ N., long. $154^{\circ} 56'$ W.; surface temperature, 74° F.

Clio (Creseis) acicula.

Clio pyramidata.

Clio (Styliola) subula.

Cavolinia inflexa.

Between Stations 264 and 265. August 24, 1875 ; Sandwich Islands to Tahiti ; lat. $13^{\circ} 15'$ N.; long. $152^{\circ} 2'$ W.; surface temperature, 78° F.

Limacina inflata.

Peraclis reticulata.

Limacina lesueuri.

Clio (Creseis) virgula.

Limacina trochiformis.

Clio (Hyaloclylix) striata.

STATION 269. September 2, 1875 ; Sandwich Islands to Tahiti ; lat. $5^{\circ} 54'$ N., long. $147^{\circ} 2'$ W.; surface temperature, $81^{\circ} \cdot 2$ F.

Cavolinia longirostris.