

STATION 122. September 10, 1873; off the coast of South America, between Pernambuco and Bahia; lat. $9^{\circ} 5' S.$, long. $34^{\circ} 50' W.$; depth, 350 fathoms; bottom, red mud.

Limacina inflata.

Limacina lesueuri.

Limacina bulimoides.

Peracelis reticulata.

Peracelis bispinosa.

Clio (Creseis) acicula.

Clio (Styliola) subula.

Clio pyramidata.

Cuvierina columnella.

Cavolinia trispinosa.

Cavolinia quadridentata.

Cavolinia longirostris.

Cavolinia uncinata.

Cavolinia inflexa.

STATION 164. June 12, 1874; off Sydney; lat. $34^{\circ} 8' S.$, long. $152^{\circ} 0' E.$; depth, 950 fathoms; bottom, green mud.

Clio (Styliola) subula.

Clio pyramidata.

Cavolinia trispinosa.

STATION 174. August 3, 1874; off Kandavu Island; lat. $19^{\circ} 6' 0'' S.$, long. $178^{\circ} 14' 20'' E.$; depth, 140 fathoms; bottom, coral mud.

Limacina inflata.

STATION 185. August 31, 1874; off Raine Island; lat. $11^{\circ} 35' 25'' S.$, long. $144^{\circ} 2' 0'' E.$; depth, 135 fathoms; bottom, coral sand.

Limacina inflata.

Limacina lesueuri.

Limacina bulimoides.

Clio (Creseis) virgula.

Clio (Creseis) acicula.

Clio (Hyalocylix) striata.

Clio (Styliola) subula.

Clio pyramidata.

Cuvierina columnella.

Cavolinia trispinosa.

Cavolinia quadridentata.

Cavolinia longirostris.

Cavolinia gibbosa.

Cavolinia uncinata.

Cavolinia inflexa.

STATION 219. March 10, 1875; Admiralty Islands to Yokohama; lat. $1^{\circ} 54' 0'' S.$, long. $146^{\circ} 39' 40'' E.$; depth, 150 fathoms; bottom, coral mud.

Limacina inflata.

Limacina trochiformis.

Limacina bulimoides.

Clio (Creseis) virgula.

Clio (Creseis) acicula.

Clio (Creseis) conica.

Clio (Hyalocylix) striata.

Clio (Styliola) subula.

Clio pyramidata.

Cavolinia trispinosa (young, as
" *Cleodora compressa* ").

Cavolinia quadridentata.

Cavolinia longirostris (young, as
" *Hyalæa lævigata* ").