"Corniger. Hands without fangs: proboscis incurved subulate: sides of the thorax with a double horn.

Inhabits the Norway Seas.

Body of 11 short segments, whitish edged with red, the 5 hind ones carinate and spinous on the back: under the thorax each side are 2 horns united at the base: tail with numerous bifid styles.

"* Linearis. Hands 4 with a single fang: legs 10.

Pall. Spicil. Zool. 9. tab. 4. fig. 15, Baster. tab. 4. fig. 2.

Inhabits the shores of Europe and America.

"* Atomos. Hands 4 with a single fang: legs 14, with two oval vesicles each side between the fourth and fifth pair.

Pennant Brit. Zool. iv. tab. 12. fig. 22.

Inhabits *Europe*, in running water, and is so very minute as to be seldom visible to the naked eye.

"* Salinus. Legs 20 spreading: tail subulate," &c. (not an Amphipod).

"* Stagnalis. Hands without fangs: legs 22: tail cylindrical bifid," &c. (not an Amphipod).

"Gibbosus. Oblong, gibbous; antennæ folded and very long.

Inhabits Portugal; small.

Body smooth yellowish speckled with brown: head thick obtuse with a large green spot: antennæ bent under the body, folded and 3 times as long as the body: tail with 3 sharp cleft leaves.

"Esca. Hands without fangs: tail jointed subulate and cleft at the tip," &c. (not an Amphipod).

"Medusarum. Hands 4 with a single fang: head very obtuse.

Stroem, Sundm. 188. tab. 1. fig. 12, 13.

Inhabits Norway, under Medusæ.

"Filiformis. Linear; legs 10, the middle ones larger.

Inhabits Melacea. Amoen. Acad. 6. p. 415. n. 99."

After "120. Monoculus" comes "121. Onisous. Jaw truncate denticulate: lip bifid: antennæ setaceous, 2-4: body oval, consisting of about 14 transverse segments: legs 14."

Under section "A. Feelers 0: antennæ often 4, sessile: Cymothoa," are given among many others:—

"Ceti. Ovate with distinct segments: third and fourth pair of legs linear and unarmed.

Seba. Mus. 1. tab. 90. fig. 5. Degeer, 7. t. 42. f. 6, 7.

Inhabits the Northern Seas, on Whales."

"Aculeatus. Thorax naked: back with 3 rows of spines.

Act. Petrop. 1778, 1. p. 247. tab. 8. fig. 1.

Inhabits the White Sea. Body carmine."

"Cuspidatus. Thorax articulate tuberculate: the 6 dorsal segments cuspidate.

Inhabits the White Sea. Act. Petrop. 1778. tab. 8. fig. 3.

Antennæ 4: tail tufted at the sides."

"Fuscus. Brown; shell carinate with a white spot on the thorax.

Inhabits Denmark. II. Mull. Zool. Dan. 2476.

"Medusarum. A little compressed: front obtuse; antennæ very short and pendant: hands 4 compressed and cut.

Stroem. Sundm. 1. p. 188. tab. 1. fig. 12, 13.

Found under the folds of the Medusa Capillata.

"Cicada. Compressed, sublinear with four spurious hands: upper antennæ shorter: tail smooth on the back.

Inhabits Greenland Seas. Fab. fn. Groen. p. 258. n. 233.