

- 3-ii-7-mi paris* latus. Long. corp. usque ad 5 mm. Colore griseo cum maculis parvis nigris. Femina feminae *C. difformi* similis."
60. *Cerapus pugnax* (Dana), is followed by a quotation of the description, and "Var. *Pontica* m. *Antennæ superiores* flagello longitudine variabili, plus minusve breviora quam pedunculum. *Pedes 2-di paris* digito minus elongato, articulo ejus 1-mo crasso, 2-do margine posteriore incrassato, apice obtuso setis nonnullis obsito. *Oculi* rubri. Long. corp. usque ad 3.8 mm. Color ut in præcedente." The quotations which I have omitted in regard to this and the preceding species are given in V. Carus, prodr. Faun. Medit., p. 393, but Czerniavski's own descriptions of his varieties are omitted by Carus.
- "61. *Cerapus bidens*, nov. sp. *Varietati ponticæ Cerapi macrodactyli* simillimus, nisi *pedes paris 2-di* digito immobili paulo curvato et in apice bidentato, articulo 5-to intus maxime eroso, tuberculo subbasali valde forti. Long. corp. usque ad 5.2 mm." These differences, from a species confessedly variable, are but of doubtful specific value. The species is not noticed by Victor Carus, Faun. Medit., except in so far as he intimates that the name is preoccupied, by giving "*C. bidens*, V. Crs. (nec Czern.)," to take the place of Costa's *Ericthonius bidens*.
- Gen. 47. *Podocerus*, Leach, is given with the synonyms, *Ischyrocerus*, Kröyer, *Cerapus* (pars), Milne-Edwards, *Jassa* Leach, Milne-Edwards, *Gammarus* (pars), Rathke, *Cratophium*, Dana. "62. *Podocerus Ocius*, Sp. Bate;" 63. *Podocerus dentex*, n. s., Tab. vi. fig. 35, is thus described:—"Mas. Frons obtusa. *Antennæ superiores* inferioribus paulo breviores et multo debiliores, flagello 3-articulato, articulo præcedente non multo longiore, articulis valde decrescentibus, filis olfactoriis tenuioribus in margine inferiore instructo, flagello secundario rudimentario; *a. inferiores* incrassatæ, flagello 3-articulato, articulis valde decrescentibus, sub apice spinis unguiformibus duabus armato. Ambo paria antennarum margine inferiore sat setoso. *Pedes paris 1-mi* manu pyriformi, triplum majore quam carpo, palma setulosa et spinulis nonnullis (3) subbasalibus armata, ungue forti, curvato, margine posteriore paulo dentato; *p. 2-di* manu magna, elongata, palma excavata, dentibus duobus posterioribus magnis et duobus vel tribus subapicalibus multo minoribus terminata setis plumosis dense obsita, ungue curvato, margine posteriore incrassato; *p. 3-ii* et *4-ti* breviores ungue minore, vix curvato. Long. corp. usque ad 4.3 mm. Color flavescens-brunescens, maculis nigris. *Oculi* nigri."
- "Gen. 48. *Sunamphithoë* (Sp. Bate), emend.," is thus defined:—
- "Ut *Amphithoë*. *Telson* crassum et vel uno hamulo vel duobus terminatum. *Pedes paris* 5-ti, 6-ti, 7-mi tarso (articulus 6-tus Sp. Bate) ad apicem dilatato." In the synonymy Czerniavski refers to *Amphithoë podoceroïdes*, Rathke, and *Amphithoë bicuspis*, Heller. He also assigns to this genus, "64. *Sunamphithoë valida* nov. sp., Tab. vi. fig. 36.," thus described:—"Mas. *Oculi* ovales. *Antennæ superiores* segmentum 6-tum thoracis attingentes, inferioribus paulo longiores, pedunculo apice articulum 3-ium inferiorum æquante, flagello filiformi duplo longiore quam pedunculus, 12-15 articulato (articulis in apice filo olfactorio instructis); *a. inferiores* validiores, pedunculo longissimo (flagello superiorum æquali), flagello brevissimo, fere triplum breviora quam pedunculus (3/4 articuli 4-ti æquante), articulis 7-9 valde decrescentibus. Ambo paria antennarum setulis paucis perbrevis instructa. *Pedes paris 1-mi* et *2-di* maxime inæquales, *p. 1-mi* carpo sat magno, triangulari, manu subpyriformi, palma obliqua, denticulo spiniformi postice terminata, ungue in margine posteriore leviter dentato; *p. 2-di* manu valida semi-ovali, palma transversa, coucava, postice dentem (fere indicem) validum, rotundatum exhibente, ungue maxime curvo, postice dentato. *Pedes ceteri* ut in *S. hamulo* Sp. Bate conformati, *p. 7-mi* tarso ad apicem maxime dilatato. *Telson* crassum, retrorsum paulo angustatum, marginibus lateralibus rectis, hamulis duobus fortibus terminatis. Flavescens, cum maculis nonnullis parvis fuscis. Femina. *Pedes paris 2-di* 1-mo non majores,