

nearly straight. Branchial vesicles pear-shaped, much broader below than above, not as long as the first joint. Marsupial plates broad and long. First joint reaching a little beyond the side-plate, with setæ and setules along the margins; third joint longer than the fourth, shorter than the fifth, a little decurrent in front; there are a few setules and spinules on the margins and apices of these joints; the finger long, thin, pointed and curved, more than half the length of the fifth joint.

*Second Peræopods.*—Side-plates rather deeper than broad. Branchial vesicles pear-shaped, bent, as long as the first joint. Marsupial plates broad, not as long as the side-plates. Neither first nor second joint reaching the end of the side-plate; the third joint longer than either the fourth or the fifth; the fourth shorter than the fifth, each with four groups of spines on the hind margin; the finger broad, shorter than in the preceding pair, much curved at the tip, with the inner margin a little serrate.

*Third Peræopods.*—Side-plates deeper than broad. Branchial vesicles short, widest just below the neck. First joint of the limb not expanded above, but forming a rounded lobe below and behind which partially overlaps the second joint, with nine spinules on the hind margin, and five or six on the lower half of the front; second joint with spines at two points in front; third joint much broader as well as longer than the fourth or fifth, with spines at six points in front, and spinules at as many on the convex decurrent hind margin; the fourth joint shorter than the fifth, each with spines at four points of the front margin; the finger much more than half the length of the fifth joint, curved at the tip, the inner margin smooth.

*Fourth Peræopods.*—Side-plates similar to the preceding pair, but smaller. First joint evenly expanded or a little wider above than below, with a few spines along the front margin, the hinder almost completely smooth, the rounded lower margin partially overlapping the second joint behind; the third and fifth joints rather larger than in the third peræopods.

*Fifth Peræopods.*—The side-plates smaller than the preceding pair. The first joint wider than in the preceding pair, and behind quite overlapping the second joint; the third joint shorter in front than in the fourth peræopods; the finger with serrate inner margin.

*Pleopods.*—Coupling-spines seemingly minute; a single cleft spine on the inner ramus; joints of the rami seven to nine in number.

*Uropods.*—Peduncles of the first pair rather longer than the rami, with spines on the pectinate upper edges; inner ramus a little shorter than the outer, with two marginal spines, the outer with three, both with the upper edges pectinate; peduncles of the second pair subequal in length to the inner ramus; the outer ramus much shorter than the inner, with one marginal spine, the inner with two, both with pectinate upper edges; peduncles of the third pair a little shorter than the ramus, with four spines along the margin; the basal portion of the ramus shorter than the nail, carrying two spines on the upper margin, the upper edge of the nail pectinate.