

*Lower Antennæ*.—Gland-cone very prominent; third (first free) joint of the peduncle curved, a very little longer than the fourth joint, which is distally a little curved in the opposite direction; the fifth joint about four-fifths the length of the fourth; the slender first joint of the flagellum considerably less than half the length of the last joint of the peduncle, the second and third joints quite minute.

*Mandibles*.—The left mandible with a very prominent tooth at the top or front of the long straight cutting edge, the secondary plate not much shorter than the principal, a minute tubercle on the surface adjoining its hind margin; the cutting edge on the right mandible more convex; the first joint of the palp very long, curved, not very broad, the second and third joints subequal in length, together much shorter than the first, the third more slender than the second.

*Maxillæ* of the usual pattern, the second pair much smaller than the first.

*Maxillipeds*.—The distal margin of the inner plate very sinuous, the outer plates broad and long.

*First Gnathopods*.—The side-plates with the lower front angle acute, with a diagonal ridge of the under surface directed to it. The first joint pretty evenly broad except at the neck, the front margin indented at the top and microscopically pectinate; the third joint widening greatly from the narrow neck, so that the distal breadth is much greater than the length of the joint, the convex hind margin pectinate on the lower half, the front forming a rounded apex, the distal margin extremely sinuous; the wrist widening immediately from the broad neck to a breadth a little less than that of the third joint, then narrowing greatly, so as to project but little behind the hand and scarcely at all in front, the sinuous hind margin longer than the front, pectinate, the hinder part of the distal margin pectinate with three or four little teeth; the hand smooth, shorter than the wrist, its length equalling or little exceeding the wrist's greatest breadth; the finger about a third of the length of the hand.

*Second Gnathopods*.—The first joint with the front margin nearly straight; the third joint a little longer than in the first gnathopods; the wrist pectinate with six little teeth on the round front apex, the triangular process behind shorter than the trunk, broad at the base, shorter than the hand, pectinate on both margins, the pectination of the hind margin continued some way up the trunk of the joint; the hand rather widened at the centre, pectinate along the hind margin; the finger scarcely more than a third of the length of the hand.

*First and Second Peræopods* as in *Parapronoë clausi*.

*Third Peræopods* like those of *Parapronoë clausi*, but the bent process within the side-plates is more narrowly produced, and the fifth joint is more decidedly longer than the fourth.

*Fourth Peræopods* very near to those of the species just mentioned; the two limbs of the specimen differ slightly in regard to the first joint, since in one the apex of the