

GEOGRAPHICAL DISTRIBUTION.

LIST OF STATIONS,

SHOWING THE PHYSICAL CONDITIONS AND THE SPECIES OF ANOMURA OBTAINED
AT EACH.

STATION VIIp. Off Tenerife, Canaries, February 10, 1873; lat. 28° 35' N., long. 16° 5' W.;
depth, 78 fathoms; bottom, volcanic sand; surface temperature, 64°. Dredged.

Pagurus calidus, Risso.

Spiropagurus elegans, Miers.

Anapagurus pusillus, n. sp.

Galathea dispersa, Spence Bate.

“Label lost, probably Madeira.” More probably Bay of Biscay, in deep water.

Eupagurus excavatus (Herbst), var. *meticulosa*, Roux.

STATION 23. Off Sombrero, West Indies, March 15, 1873; lat. 18° 24' N., long. 63° 28'
W.; depth, 450 fathoms; bottom, Pteropod ooze; surface temperature, 76°. Dredged.

Munida sp.

Munidopsis sigsbei (A. Milne-Edwards).

Uroptychus nitidus (A. Milne-Edwards).

Elasmonotus armatus, A. Milne-Edwards).

St. Thomas, West Indies, shallow water.

Hypoconcha sabulosa (Herbst).

Porcellana sayana (Leach).

STATION 24. Off Culebra Island, West Indies, March 25, 1873; lat. 18° 38' 30" N.,
long. 5° 55' 30" W.; depth, 390 fathoms; bottom, Pteropod ooze; surface tempera-
ture, 76°. Dredged.

Porcellana robertsoni, n. sp.

Munida microphthalma, A. Milne-
Edwards.

Elasmonotus armatus, A. Milne-Edwards.

Uroptychus nitidus (A. Milne-Edwards).