

Bermuda, on the sandy shore and in shallow water.

Remipes scutellatus (Fabricius). | *Petrolisthes armatus* (Gibbes).

STATION 49. South of Halifax, Nova Scotia, May 20, 1873; lat. 43° 3' N., long. 63° 39' W.; depth, 85 fathoms; bottom, gravel, stones; surface temperature, 40°·5; bottom temperature, 35°. Dredged.

Eupagurus pubescens (Kröyer), var. *kroyeri*, Stimpson.

STATION 56. Off Bermuda, May 29, 1873; lat. 32° 8' 45" N., long. 64° 54' 35" W.; depth, 1075 fathoms; bottom, coral mud; bottom temperature, 38°·2; surface temperature, 72°·5. Dredged.

Parapagurus abyssorum, A. Milne-Edwards. | *Munidopsis serratifrons* (A. Milne-Edwards).

STATION 68. Between Bermuda and the Azores, June 24, 1873; lat. 38° 4' N., long. 39° 19' W.; depth, 2175 fathoms; bottom, Globigerina ooze; bottom temperature, 36°·2; surface temperature, 70°. Trawled.

Parapagurus abyssorum, A. Milne-Edwards, var. *scabra*, nov.

STATION 73. West of the Azores, June 30, 1873; lat. 38° 30' N., long. 31° 14' W.; depth, 1000 fathoms; bottom, Pteropod ooze; bottom temperature, 39°·4; surface temperature, 69°. Dredged.

Pagurodes (?) sp.

STATION 75. Off Fayal, Azores, July 2, 1873; lat. 38° 38' 0" N., long. 28° 28' 30" W., depth, 50–90 fathoms; bottom, volcanic mud; surface temperature, 70°·2. Dredged.

Anapagurus pusillus, n. sp. | *Galathea* sp.

St. Vincent, Cape Verde Islands.

Remipes scutellatus (Fabricius). | *Galathea* sp.
Pagurus callidus, Risso. | *Pachycheles barbatus*, A. Milne-Edwards.
Eupagurus excavatus (Herbst), var. |
meticulosa, Roux.

STATION 78. South of San Miguel, Azores, July 10, 1873; lat. 37° 26' N., long. 25° 13' W.; depth, 1000 fathoms; bottom, volcanic mud; surface temperature, 71°. Dredged.

Lithodes agassizii, S. I. Smith.