

Reefs at Levuka, Fiji Islands.

Catapagurus australis, n. sp.

Kandavu, Fiji Islands, from the shore.

Cænobia rugosa, H. Milne-Edwards. | *Cænobia perlata*, H. Milne-Edwards.

STATION 174D. Off Ngaloa, Kandavu, Fiji Islands, August 3, 1874 ; lat. $19^{\circ} 5' 50''$ S., long. $178^{\circ} 16'' 20''$ E. ; depth, 210 fathoms ; bottom, coral mud ; surface temperature, $77^{\circ}.7$. Dredged.

Lyreidus tridentatus, De Haan. | *Munida spinicordata*, Henderson.

Api, New Hebrides, from the shore.

Cænobia rugosa, H. Milne-Edwards. | *Cænobia perlata*, H. Milne-Edwards.

STATION 186. Flinder's Passage, Cape York, North Australia, September 8, 1874 ; lat. $10^{\circ} 30'$ S., long. $142^{\circ} 18'$ E. ; depth, 8 fathoms ; bottom, coral mud ; surface temperature, $77^{\circ}.2$. Dredged.

Mastigochirus quadrilobatus, Miers. | *Petrolisthes annulipes* (White) Miers.

Pagurus imbricatus, H. Milne-Edwards. | *Pachycheles pulchellus* (Haswell).

Polyonyx obesulus (White) Miers.

Torres Strait.

Spiropagurus spiriger (De Haan).

STATION 187. Cape York, North Australia, September 9, 1874 ; lat. $10^{\circ} 36'$ S., long. $141^{\circ} 55'$ E. ; depth, 6 fathoms ; bottom, coral mud ; surface temperature, $77^{\circ}.7$.

Diogenes guttatus, n. sp.

STATION 188. Arafura Sea, south of Papua, September 10, 1874 ; lat. $9^{\circ} 59'$ S., long. $139^{\circ} 42'$ E. ; depth, 28 fathoms ; bottom, green mud ; surface temperature, $78^{\circ}.5$. Dredged and trawled.

Spiropagurus spiriger (De Haan). | *Catapagurus australis*, n. sp.

Pachycheles pulchellus (Haswell).

STATION 190. Arafura Sea, south of Papua, September 12, 1874 ; lat. $8^{\circ} 56'$ S., long. $136^{\circ} 5'$ E. ; depth, 49 fathoms ; bottom, green mud ; surface temperature, $79^{\circ}.2$. Trawled.

Spiropagurus spiriger (De Haan). | *Galathea australiensis*, Stimpson.