

STATION 218. North of Papua, March 1, 1875; lat. $2^{\circ} 33' S.$, long. $144^{\circ} 4' E.$; depth, 1070 fathoms; bottom, blue mud; bottom temperature, $36^{\circ} \cdot 4$; surface temperature, 84° . Trawled.

Parapagurus abyssorum, A. Milne-Edwards.

Munidopsis brevimana, Henderson.
Elasmonotus latifrons, Henderson.

STATION 219. Off Nares Harbour, Admiralty Islands, March 10, 1875; lat. $1^{\circ} 54' 0'' S.$, long. $146^{\circ} 39' 40'' E.$; depth, 150 fathoms; bottom, coral mud; surface temperature, 84° . Trawled.

Munida squamosa, Henderson.
Munida proxima, Henderson.

Munida inornata, Henderson.
Elasmonotus lævigatus, Henderson.

Wild Island, Admiralty Islands.

Cænobita clypeata (Herbst).

Cænobita rugosa, H. Milne-Edwards.

Tracy Island, Nares Harbour, Admiralty Islands.

Cænobita rugosa, H. Milne-Edwards.

Admiralty Islands; depth, 16 to 25 fathoms.

Pagurus dearmatus, n. sp.

Spiropagurus spiriger (De Haan).

Off Yokoska, Japan; depth, 5 to 20 fathoms.

Cryptodromia japonica, n. sp.

Off Yokohama, Japan.

Eupagurus constans, Stimpson.

STATION 237. South of Japan, June 17, 1875; lat. $34^{\circ} 37' N.$, long. $140^{\circ} 32' E.$; depth, 1875 fathoms; bottom, blue mud; bottom temperature, $35^{\circ} \cdot 3$; surface temperature, 73° . Trawled.

Parapagurus abyssorum, A. Milne-Edwards.

Munidopsis subsquamosa, Henderson.

Tahiti, Society Islands, near the reefs, September 28, 1875.

Cænobita rugosa, H. Milne-Edwards.

Calcinus tibicen (Herbst).

Reefs at Papiete, Society Islands.

Pagurus deformis, H. Milne-Edwards.