

fauna of the Cape Verde Islands (which I include in this section) shows a decided Lusitanian facies.

Remipes scutellatus (Fabricius), II.
Pagurus calidus, Risso, III.
Eupagurus excavatus (Herbst), var.
meticulosa, Roux, III.

Pachycheles barbatus, A. Milne-Edwards.
Munida microphthalma, A. Milne-Edwards, II., X.

V. THE SOUTH AFRICAN PROVINCE.

Includes the south-west coast of Africa from the Tropic of Capricorn to Cape Agulhas, and the Tristan da Cunha group.

Dromidia spongiosa, Stimpson.
 „ *bicornis*, Studer.
Eudromia frontalis, n. gen. et sp.
Pseudodromia latens, Stimpson.
Diogenes brevirostris, Stimpson.
Pagurus granulatus, Olivier.
Eupagurus tristanensis, n. sp.
Anapagurus pusillus, n. sp., III.

Parapagurus dimorphus (Studer), XII.
 * „ *abyssorum*, A. Milne-Edwards, II., III., VI., VII., XI.
Porcellana streptocheles, Stimpson.
Galathea labidolepta, Stimpson (?).
 „ sp.

THE INDO-PACIFIC REGION.

VI. THE INDO-MALAYAN PROVINCE.

This province is one of considerable extent. It includes the Malay Archipelago, the north coast of Australia, and extends in a northerly direction so as to take in the greater part of the Chinese seas, while it also comprises the coasts of India and Ceylon.

Homola orientalis, n. sp.
 * *Homologenus* (?) sp.
Latreillopsis bispinosa, n. gen. et sp.
Latreillia valida, De Haan.
Raninoides personatus (White), Henderson.
Notopoides latus, n. gen. et sp.
Notopus ovalis, n. sp.
Cosmonotus grayii, Adams and White.
Remipes testudinarius, Latreille.

Mastigochirus quadrilobatus, Miers.
Albunea microps (White), Miers.
Birgus latro (Linné).
Canobita clypeata (Herbst).
 „ *rugosa*, H. Milne-Edwards, X.
Diogenes guttatus, n. sp.
Pagurus striatus, Latreille.
 „ *imbricatus*, H. Milne-Edwards.
 „ *dearmatus*, n. sp.