

<i>Galathea aculeata</i> , Haswell, VI.	<i>Elasmonotus miersii</i> , Henderson.
* <i>Munida microphthalma</i> , A. Milne-Edwards, II., IV.	„ <i>debilis</i> , Henderson.
„ <i>normani</i> , Henderson.	* <i>Uroptychus spinimarginatus</i> , Henderson, VI.
„ <i>granulata</i> , Henderson.	* „ <i>australis</i> , Henderson, VI., VIII.
„ <i>militaris</i> , Henderson, VI.	„ <i>politus</i> , Henderson.
„ <i>tuberculata</i> , Henderson.	
„ <i>spinicordata</i> , Henderson.	

## XI. THE PERUVIAN PROVINCE.

This includes the coasts of Chili and Peru; possibly it extends further north than the boundary of the latter country.

* <i>Parapagurus abyssorum</i> , A. Milne-Edwards, II., III., V., VI., VII.	* <i>Munidopsis antonii</i> (A. Milne-Edwards), XII.
<i>Petrolisthes violaceus</i> (Guérin).	* <i>Galacantha bellis</i> , n. sp.
„ <i>validus</i> (Dana), XII.	

## THE ANTARCTIC REGION.

## XII.

As already indicated, this region is held to include all these species taken to the south of 40° S. latitude. I have, however, comprised certain Antarctic forms taken at Station 320, off the Rio de la Plata.

<i>Lithodes murrayi</i> , n. sp.	* <i>Munida spinosa</i> , Henderson.
<i>Paralomis verrucosus</i> (Dana).	<i>Munidopsis antonii</i> (A. Milne-Edwards), XI.
„ <i>aculeatus</i> , n. sp.	* „ <i>subsquamosa</i> , Henderson, var. <i>aculeata</i> , nov.
* „ <i>formosus</i> , n. sp.	„ <i>trifida</i> , Henderson.
<i>Eupagurus comptus</i> (White), var. <i>jugosa</i> , nov.	<i>Elasmonotus asper</i> , Henderson.
<i>Parapagurus dimorphus</i> (Studer), V.	<i>Ptychogaster milne-edwardsi</i> , Henderson.
* <i>Pagurodes inarmatus</i> , n. sp., IX.	<i>Uroptychus insignis</i> , Henderson.
<i>Petrolisthes validus</i> (Dana), XI.	„ <i>parvulus</i> , Henderson.
<i>Porcellanella triloba</i> , White, VI.	
<i>Munida subrugosa</i> (White).	