E	rvilia castanea (Montagu). Obtained also at Stations 75 and 78, 450 and 1000 fathoms.	STATION VIIP.
P	sammobia costulata, Turton. Obtained at no other locality by the Challenger.	
	Recorded from North Atlantic and Mediterranean.	
V_{i}	enerupis irus (Linné). Obtained at no other locality by the Challenger. Recorded	
	from North Atlantic.	
V	enus (Ventricola) casina, Linné. Obtained also at Station 75, 450 fathoms.	
	Recorded from North Atlantic and Mediterranean.	
	" (") effossa, Bivona. Obtained also at Station 75, 450 fathoms.	
C_{i}	ytherea (Callista) chione, Linné. Obtained at no other locality by the Challenger.	
	Recorded from North Atlantic and Mediterranean. Fossil-	
	Coralline Crag and Tertiary formations of Italy and Sicily.	
C_{i}	firce minima (Montagu). Obtained also at Station 75, 450 fathoms. Recorded	
	from North Atlantic and Mediterranean.	
C c	ardium (Acanthocardium) papillosum, Poli. Obtained also at Station 75, 50 to	
	450 fathoms.	
	" (Papyridea) transversale, Deshayes. Obtained also at Station 75,	
17.020	450 fathoms. Recorded from Alboran Island.	
C_{i}	hama gryphoides, Linné. Obtained also at Station 75, 450 fathoms. Recorded	
_	from North Atlantic and Mediterranean.	
D	hiplodonta apicalis, Philippi. Obtained at no other locality by the Challenger.	
	Recorded from Canaries.	
A_{\cdot}	starte macandrewi, Smith. Obtained at no other locality by the Challenger.	
C	Recorded from Canaries.	
	ardita calyculata (Linné). Obtained also at Station 162, 38 to 40 fathoms. Recorded from Mediterranean and Fiji.	
P	ectunculus stellatus (Bruguière). Obtained at no other locality by the Challenger.	
	imopsis minuta (Philippi). Obtained also at Stations II., VIII., 24, and 75.	
	ima squamosa, Lamarck. Obtained also at Station 212, 10 fathoms. A widely-	
13	distributed species.	
P	Pecten pusio (Linné). Obtained also at Station 75, 450 fathoms.	
	, corallinoides, d'Orbigny. Obtained also at Cape Verdes. Recorded	
	from Canaries.	
GASTEI	ROPODA (Watson, Zool. pt. 42).	
	rochus (Ziziphinus) striatus, Linné. Obtained also at Stations 75 and 122,	
1	450 and 350 fathoms. Recorded from North Atlantic	
	and Mediterranean. Fossil-Upper Miocene all over	
	Europe onwards.	