## SUMMARY OF RESULTS.

Cladocora arbuscula, M.-Edwards and Haime. Dead fragment; obtained also in STATION 33. Simon's Bay, Cape, 10 to 20 fathoms.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiomusium cancellatum, n.sp. (?). Obtained also at Station 236, 420 to 775 fathoms.

Ophiopyren longispinus, n.g., n.sp. Obtained also at Stations 23 and 24.

Amphilepis norvegica, Ljungman (?). Obtained also at Stations 45 and 46, 1240 and 1350 fathoms.

Ophiacantha troscheli, n.sp. Obtained at no other locality by the Challenger. Recorded subsequently from North Atlantic ("Blake").

Ophiomitra chelys (Wyville Thomson), n.sp. (var. ?). For distribution see Station 3. Astroschema brachiatum, n.sp. Obtained at no other locality.

ANNELIDA (M'Intosh, Zool. pt. 34).

Eulepis wyvillei, n.sp. One specimen; obtained at no other locality.

Eunotomastus grubei, n.g., n.sp. One fragmentary specimen ; obtained at no other locality. Only species of the genus.

Placostegus assimilis, n.sp. One specimen; obtained at no other locality.

Spirobranchus occidentalis, n.sp. One specimen, attached to tube of Placostegus assimilis; obtained at no other locality.

OSTRACODA (Brady, Zool. pt. 3).

"

- Pontocypris trigonella, Sars. Obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil— Post-Tertiary of Scotland.
- Bairdia foveolata, Brady (?). Obtained also at Stations 93, 94, 162, 187, 189, 279, Hong Kong Harbour, and Admiralty Islands, 6 to 1150 fathoms.
- Cythere bermudæ, Brady. Obtained at no other locality by the Challenger. Recorded from Colon-Aspinwall.
  - " fungoides (Brady). Obtained also at Stations 187 and 189, 6 to 8 and 28 fathoms. Recorded from Australia.

Xestoleberis curta, Brady. Obtained also at Stations 149, 187, 300, Port Jackson, and Sandwich Islands, 2 to 1375 fathoms. Recorded from West Indies.

Asterope sp. (?). A few imperfect specimens.

Cytherella pulchra, Brady. Obtained also at Stations 167, 344, and Port Jackson, 2 to 420 fathoms. Recorded from Australia.

irregularis, n.sp. A few detached valves; obtained at no other locality.