STATION 75. CORALS (Moseley, Zool. pt. 7).

Caryophyllia clavus, Scacchi, var. smithi, Duncan (?). Several small specimens; obtained also at Station 308, 175 fathoms.

Paracyathus de filippii (Duchassaing and Michelotti). Numerous specimens (50 fathoms); obtained also at Station 190, 49 fathoms.

HYDROIDA (Allman, Zool. pts. 20 and 70).

Polyplumaria pumila, n.sp. Several fragmentary specimens; obtained at no other locality.

Aglaophenia filicula, n.sp. Obtained at no other locality.

acacia, n.sp. Obtained at no other locality.

Halecium beanii, Johnston. Obtained also at Station 163A, 150 fathoms. Recorded from Britain.

Perisiphonia filicula, n.g., n.sp. Obtained also at Station 163A, 150 fathoms.

The only other species of the genus (Perisiphonia pectinata) was taken at Station 169.

Sertularia laxa [exigua], n.sp. Obtained at no other locality.

Diphasia pinaster (Ellis and Solander). Obtained at no other locality by the Challenger. Recorded from European seas.

Thuiaria pharmacopola, n.sp. Obtained at no other locality.

ASTEROIDEA (Sladen, Zool. pt. 51).

Astropecten hermatophilus, n.sp. Obtained at no other locality.

Chætaster longipes (Retzius), Sars. Obtained also at Station 36.

Ophidiaster attenuatus, Gray. Obtained at no other locality by the Challenger. Recorded from Mediterranean.

ophidianus (Lamarck), Agassiz (?). One small specimen; obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean.

OPHIUROIDEA (Lyman, Zool. pt. 14).

Ophiothrix lütkeni, Wyville Thomson (?). Obtained at no other locality by the Challenger.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Spatangus purpureus, Müller. (50 to 90 fathoms); obtained also at Bermuda, 100 fathoms.

Brissus damesi, n.sp. Obtained also at Station 122, 350 fathoms.

ANNELIDA (M'Intosh, Zool. pt. 34).

Lepidonotus squamatus (Linné). One specimen; obtained at no other locality by the Challenger. Recorded from Virginian coast.