Stomatella (Gena) nigra, Quoy and Gaimard (?). One specimen; obtained also at Port Jackson, 2 to 10 fathoms. Recorded from Tongatabu, Sandwich Islands, and West Africa.

Fernando Noronha.

Phasianella sp. (?).

Nerita ascensionis, Gmelin. (St. Michael's Rocks); obtained at no other locality by the Challenger. Recorded from Ascension.

Solarium sp. (?).

Scalaria (Cirsotrema) hellenica, Forbes. Obtained at no other locality by the Challenger. Recorded from North Atlantic and Mediterranean. Fossil — Austrian Miocene and Piedmontese Pliocene.

Nassa (Hima) capillaris, n.sp. Obtained at no other locality.

Oliva (Olivella) fulgida, Reeve. Obtained at no other locality by the Challenger. Recorded from West Indies.

,, (,,) pulchella, Duclos (?). Two fragments; obtained at no other locality by the Challenger. Recorded from West Indies.

Columbella mercatoria (Linné). Obtained also at the Cape. Recorded from North Atlantic.

sp. (?).

Marginella (Persicula) sagittata, Hinds. Obtained at no other locality by the Challenger. Recorded from Brazil, Bahamas, Caribbean Sea, and Australia.

Cypræa spurca, Linné. Obtained also at Stations VIIP. and 122.

Mitrularia uncinata (Reeve) (?). Obtained also at Station 122, 350 fathoms, and
North Atlantic, deep water. Recorded from West
Indies and Philippines.

Xenophora corrugata (Reeve). Obtained at no other locality by the Challenger. Recorded from Indian Ocean, Japan, and China seas.

Cerithiopsis sp. (?).

Litorina (Hamus) nodulosa, d'Orbigny. (St. Michael's Rocks); obtained at no other locality by the Challenger. Recorded from West Indies.

Fossarus ambiguus (Linné). Obtained also at Station 122, 350 fathoms. Recorded from North Atlantic and Mediterranean. Fossil—Post-Tertiary of Sicily.

Rissoa sp. (?).

Utriculus (Tornatina) canaliculatus (Say). Obtained at no other locality by the Challenger. Recorded from United States coast.

Cylichna noronyensis, n.sp. Obtained at no other locality. Vanikoro (?) sp.