

STATIONS 122 TO
122c.

- Cypræa cinera*, Gmelin. (350 to 400 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.
- „ *spurca*, Linné. (350 fathoms); obtained also at Station VIIp. and Fernando Noronha.
- Velutina* sp. (?). (350 fathoms).
- Natica* sp. (?). Same as at Stations 24, 85, and 120.
- Mitrularia uncinata* (Reeve). (350 fathoms); obtained also at Fernando Noronha and North Atlantic, deep water.
- Xenophora caribæa*, Petit. (350 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.
- Siliquaria (Tenagodus)* sp. (?). (350 fathoms).
- Turritella exoleta* (Linné). (350 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.
- Odostomia turrita*, Hanley (?). One specimen (350 fathoms); obtained also at Station VIIp.
- „ *unidentata* (Montague). One injured specimen (350 fathoms); obtained at no other locality by the Challenger. Recorded from Arctic, North Atlantic, and Mediterranean. Fossil—Middle Pliocene of Calabria, English Crag, glacial beds of Norway and Scotland.
- „ (*Turbonilla*) sp. (?). (350 fathoms).
- Eulimella rudis*, n.sp. (350 fathoms); obtained at no other locality.
- Aclis hyalina*, n.sp. (350 fathoms); obtained at no other locality.
- „ *sarissa*, n.sp. (350 fathoms); obtained at no other locality.
- Eulima ephamilla*, n.sp. (350 fathoms); obtained at no other locality.
- „ *sarissa*, n.sp. (350 fathoms); obtained at no other locality.
- „ *hebes*, n.sp. (350 fathoms); obtained at no other locality.
- „ sp. (?). (350 fathoms).
- Bittium* sp. (?). (350 fathoms).
- Triforis perversa* (Linné). (350 fathoms); obtained also at Stations VIIp. and 75.
- Cithna tenella* (Jeffreys). (350 fathoms); for distribution see Station 75.
- Fossarus ambiguus* (Linné). (350 fathoms); obtained also at Fernando Noronha.
- Rissoa (Alvania) deliciosa*, Jeffreys. (350 fathoms); obtained also at Station 85.
- „ *xanthias*, n.sp. (350 fathoms); obtained also at Station 24.
- „ *pernambucensis*, n.sp. (350 fathoms); obtained at no other locality.
- „ (*Ceratia*) *pachia*, n.sp. (350 fathoms); obtained at no other locality.
- „ (*Cingula*) *rustica*, n.sp. (350 fathoms); obtained at no other locality.
- Rissoina dubiosa* (Adams) (?). (350 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.