

Ophiomyxa flaccida (Say). (7 to 20 fathoms); obtained also at Station 36.

Astrophyton costosum, Seba. (7 to 20 fathoms); obtained at no other locality by the Challenger.

BAHIA.

ECHINOIDEA (Agassiz, Zool. pt. 9).

Cidaris tribuloides (Lamarck). (7 to 20 fathoms); obtained also at Cape Verdes and Fernando Noronha.

Clypeaster subdepressus (Gray). Obtained at no other locality by the Challenger.

Mellita sexforis, Agassiz. Obtained at no other locality by the Challenger.

Encope emarginata (Leske). (20 to 70 fathoms?); obtained at no other locality by the Challenger.

Metalia pectoralis, Agassiz. Obtained at no other locality by the Challenger.

HOLOTHURIOIDEA (Théel, Zool. pt. 39).

Colochirus pygmæus, n.sp. One specimen (7 to 20 fathoms); obtained at no other locality.

Thyone pervicax, n.sp. One specimen (7 to 20 fathoms); obtained at no other locality.

GEPHYREA (Selenka, Zool. pt. 36).

Thalassema baronii, Greeff. One specimen (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from Canaries.

MYZOSTOMIDA (Graff, Zool. pt. 27).

Myzostoma gigas, Lütken. (On *Antedon carinata*).

ISOPODA.

One specimen, genus and species undetermined.

MACRURA (Spence Bate, Zool. pt. 52).

Sicyonia carinata (Olivier). One specimen (20 fathoms); obtained also at St. Thomas.

Alpheus intrinsecus, n.sp. One specimen (7 to 20 fathoms); obtained at no other locality.

„ *minus*, Say. Three specimens; obtained also at St. Paul's Rocks and Fernando Noronha.

ANOMURA (Henderson, Zool. pt. 69).

Dromidia antillensis, Stimpson. Several specimens (7 to 20 fathoms); obtained at no other locality by the Challenger.