Hypoconcha panamensis, Smith (?). One specimen (shallow water); obtained at no other locality by the Challenger.

Zanclifer caribensis (Freminville). One specimen (7 to 20 fathoms); obtained at no other locality by the Challenger. Recorded from West Indies.

Petrolisthes servatus, n.sp. One specimen (7 to 20 fathoms); obtained at no other locality.

sp. (?). One specimen (shallow water).

BRACHYURA (Miers, Zool. pt. 49).

,,

Leptopodia sagittaria (Fabricius). Several specimens (7 to 20 fathoms); obtained also at Station 122, Madeira, and Cape Verdes.

Metoporaphis forficulatus, M.-Edwards. One specimen (7 to 20 fathoms); obtained also at Station 122.

Notolopas brasiliensis, n.sp. Three specimens (7 to 20 fathoms); obtained at no other locality.

Picroceroides tubularis, n.g., n.sp. Two specimens; obtained also at Fernando Noronha.

Macrocæloma trispinosa (Latreille). Four specimens (shallow water); obtained also at Bermuda.

- ", septemspinosa (Stimpson). One specimen (shallow water); obtained at no other locality by the Challenger. Recorded from Tortugas.
- " concava, n.sp. One specimen (shallow water); obtained also at Fernando Noronha.
- Mithrax cornutus, Saussure. Four specimens (shallow water); obtained at no other locality by the Challenger. Recorded from West Indies.
 - " forceps (M.-Edwards). One specimen (shallow water); obtained also at Bermuda.
- Lambrus guérinii, Capello, var. (?). Two specimens (shallow water); obtained at no other locality by the Challenger. Recorded from West Atlantic and Mauritius.
 - " serratus, M.-Edwards. Two specimens (shallow water); obtained at no other locality by the Challenger. Recorded from West Indies.

Heterocrypta granulata (Gibbes). One specimen (shallow water); obtained at no other locality by the Challenger. Recorded from North-West Atlantic.

376