

STATIONS 144A
TO 145A.

OSTRACODA (Brady, Zool. pt. 3).

- Macrocypris maculata*, Brady. (50 to 150 fathoms); obtained also at Stations 149, 162, Simon's Bay, Cape, and Amboina.
- Bythocypris reniformis*, n.g., n.sp. (50 to 150 fathoms); for distribution see Station 24.
- Bairdia villosa*, n.sp. (50 to 150 fathoms); obtained also at Stations 135, 149, and 162, 38 to 150 fathoms.
- Cythere securifer*, n.sp. Numerous specimens (50 to 150 fathoms); obtained at no other locality.
- „ *kerguelenensis*, n.sp. (50 to 150 fathoms); obtained also at Stations 149, 162, and Port Jackson, 2 to 50 fathoms.
- „ *subrufa*, n.sp. (50 to 150 fathoms); obtained also at Station 149, 20 to 50 fathoms.
- „ *parallelogramma*, n.sp. (50 to 150 fathoms); obtained at no other locality.
- „ *polytrema*, Brady. A few detached valves (depth not given); obtained at no other locality by the Challenger. Fossil—Antwerp Crag.
- „ *suhmi*, n.sp. (50 to 150 fathoms); obtained also at Station 241, 2300 fathoms.
- Krithe producta*, n.sp. (50 to 150 fathoms); widely distributed (see Station 70).
- Xestoleberis setigera*, n.sp. (50 to 150 fathoms); obtained also at Stations 149 and 151, 120 and 75 fathoms.
- Pseudocythere caudata*, Sars. (50 to 150 fathoms); obtained also at Stations 149 and 323, 20 to 1900 fathoms.

AMPHIPODA (Stebbing, Zool. pt. 67).

- Acontiostoma marionis*, n.g., n.sp. One specimen (50 to 75 fathoms); obtained at no other locality.
- Amphilocheus marionis*, n.sp. One specimen (100 fathoms); obtained at no other locality.
- Atylopsis emarginatus*, n.g., n.sp. Two specimens (310 fathoms); obtained at no other locality.
- Pardalisca marionis*, n.sp. One specimen (100 fathoms); obtained at no other locality.

ISOPODA (Beddard, Zool. pts. 33 and 48).

- Serolis septemcarinata*, Miers. Many specimens with young in the pouches (50 and 140 fathoms); obtained also at Station 149, 25 to 60 fathoms. Recorded from Crozets and Kerguelen.