### STATION 146. OSTRACODA (Brady, Zool. pt. 3).

Cythere viminea, n.sp. Single valve; obtained at no other locality.

- " dictyon, n.sp. Widely distributed (see Station 24).
- " acanthoderma, n.sp. Obtained also at Stations 64, 73, 191, 246, 296, and 302.
- " dasyderma, n.sp. Widely distributed (see Station 5).

Krithe producta, n.sp. Widely distributed (see Station 70).

## CIRRIPEDIA (Hoek, Zool. pt. 25).

Scalpellum brevicarinatum, n.sp. One specimen; obtained also at Station 147, 1600 fathoms.

- ,, tenue, n.sp. One specimen; obtained at no other locality.
- " flavum, n.sp. Three specimens; obtained at no other locality.

### AMPHIPODA (Stebbing, Zool. pt. 67).

Andania gigantea, n.sp. One specimen; obtained also at Station 147, 1600 fathoms.

# ISOPODA (Beddard, Zool. pts. 33 and 48).

Serolis antarctica, n.sp. Obtained also at Stations 122 and 147.

Eurycope sarsii, n.sp. Two specimens; obtained also at Station 147, 1600 fathoms. Arcturus spinosus, n.sp. Eleven specimens; obtained at no other locality.

### PHYLLOCARIDA (Sars, Zool. pt. 56).

Nebaliopsis typica, n.g., n.sp. One fragmentary specimen; obtained also at Station 289, 2550 fathoms.

### Schizopoda (Sars, Zool. pt. 37).

Eucopia australis, Dana. For distribution see Station 50.

# MACRURA (Spence Bate, Zool. pt. 52).

Petalidium foliaceum, n.g., n.sp. Four specimens; obtained also at Station 159, 2150 fathoms. Only species of the genus.

Glyphocrangon podager, n.sp. One specimen; obtained at no other locality.

Nematocarcinus proximatus, n.sp. Two specimens; obtained also at Stations 188,

237, 300, and 302, 28 to 1875 fathoms.

# Anomura (Henderson, Zool. pt. 69).

Pagurodes inarmatus, n.g., n.sp. Three specimens; obtained also at Station 168, 1100 fathoms.