STATION 147.

Eurycope fragilis, n.sp. One specimen; obtained also at Stations 152, 158, and 237, 1260 to 1875 fathoms.

" sp. (?) (Eurycope atlantica, n.sp.?) One fragmentary specimen. Arcturus brunneus, n.sp. Four specimens; obtained at no other locality.

SCHIZOPODA (Sars, Zool. pt. 37).

Boreomysis scyphops, Sars. Several specimens; obtained also at Stations 157 and 158, 1950 and 1800 fathoms. Recorded from North Atlantic and Arctic.

Amblyops crozetii, Willemoes-Suhm, MS., n.sp. One specimen; obtained at no other locality.

MACRURA (Spence Bate, Zool. pt. 52).

Hymenodora duplex, n.sp. One specimen; obtained at no other locality.

Pycnogonida (Hoek, Zool. pt. 10).

Nymphon hamatum, n.sp. Obtained also at Station 146.

Colossendeis gigas, n.sp. Obtained also at Stations 146 and 300.

- " leptorhynchus, n.sp. Obtained also at Stations 146, 300, and 310.
- " gracilis, n.sp. One or two specimens; obtained also at Station 146.

Phoxichilidium pilosum, n.sp. One or two specimens; obtained also at Station 157, 1950 fathoms.

GASTEROPODA (Watson, Zool. pt. 42).

Fusus (Neptunea) calathiscus, n.sp. Obtained at no other locality.

,, (,,) setosus, n.sp. Obtained also at Station 146.

Guivillea alabastrina, n.g., n.sp. One specimen; obtained at no other locality.
Only species of the genus.

Pleurotoma (Pleurotomella) papyracea, n.sp. Obtained at no other locality.

CEPHALOPODA (Hoyle, Zool. pt. 44).

Bathyteuthis abyssicola, n.g., n.sp. One specimen; obtained at no other locality.

Polyzoa (Busk, Zool. pts. 30 and 50).

Bicellaria infundibulata, n.sp. Obtained also at Station 156, 1975 fathoms.

Bugula reticulata, n.sp. Obtained also at Stations 299, 303, and 320, 600 to 2160 fathoms.

Foveolaria orbicularis, n.g., n.sp. Obtained at no other locality.

Idmonea marionensis, Busk. Obtained also at Stations 145, 151, and 320.