TONGATABU.

Natica (Mamma) aurantia, Lamarck. Obtained at no other locality by the Challenger. Recorded from Australia and Red Sea. Fossil—shores of Red Sea.

Amalthea australis, Lamarck. Obtained at no other locality by the Challenger. Recorded from Australia and Philippines.

Cæcum sepimentum, var. arcuata, de Folin. One specimen; obtained also at Tahiti, 20 fathoms. The species obtained also at Sandwich Islands, 40 fathoms. Recorded from Mauritius.

, exile, n.sp. One specimen; obtained at no other locality.

" , two other species undetermined.

Eulima sp. (?).

Mucronalia xanthias, n.sp. Obtained also at Stations 186 and 212, 8 to 20 fathoms.

Cerithium (Vertagus) asperum (Linné). Obtained also at Fiji. Recorded from

Indian and Pacific Oceans. Fossil—shores of

Red Sea.

" ( ", ), n.sp. (?).

Bittium perparvulum, n.sp. Obtained also at Station 186, Torres Strait, and Sandwich Islands, 7 to 40 fathoms.

Rissoa sp. (?).

" (?) sp.

Scaliola bella, Adams. Obtained also at Stations 185, 186, 187, Torres Strait, and Sandwich Islands, 3 to 155 fathoms. Recorded from Japan.

Actwon flammeus (Gmelin). Obtained also at Station 188 and Fiji, 28 and 12 fathoms. Recorded from Indian and Pacific Oceans.

NUDIBRANCHIATA (Bergh, Zool. pt. 26).

Platydoris eurychlamys, Bergh. One specimen; obtained also at Philippines, 10 fathoms. Recorded from Philippines.

CEPHALOPODA (Hoyle, Zool. pt. 44).

Octopus tonganus, n.sp. Three specimens; obtained at no other locality.

POLYZOA (Busk, Zool. pts. 30 and 50).

Membranipora albida, Hincks (?). Obtained also at Station 75.

Retepora apiculata, n.sp. Obtained at no other locality.

producta, n.sp. Obtained also at Philippines, 10 fathoms.

Mucronella quadrata, n.sp. Obtained at no other locality.

Idmonea radians (Lamarck). Obtained also at Sandwich Islands, 20 to 40 fathoms.

Recorded from Australian seas, New Zealand, &c.