CAPE YORK:

Tænia trichoglossi, n.sp. (?). Nine fragments (from intestine of Trichoglossus swainsoni, Cape York).

Annelida (M'Intosh, Zool. pt. 34).

Lepidonotus cristatus, Grube. One specimen (Station 186, 8 fathoms); obtained at no other locality by the Challenger. Recorded from Philippines.

Eupompe australiensis, n.sp. One specimen (Station 186, 8 fathoms); obtained at no other locality.

Eunice torresiensis, n.sp. Numerous specimens (Station 186, 8 fathoms); obtained at no other locality.

Hyalinæcia tubicola, Müller, var. papuensis, nov. (Station 186, 8 fathoms); obtained at no other locality. A widely-distributed species.

Thelepus sp. (?). Fragment (Station 186, 8 fathoms).

Myzostomida (Graff, Zool. pts. 27 and 61).

Myzostoma horologium, n.sp. (Stations 186 and 187, host uncertain).

- dentatum, n.sp. (Station 186, on Antedon variipinna).
- " intermedium, n.sp. (Station 187, on Antedon multiradiata).
- " quadrifilum, n.sp. (Station 186, on Antedon variipinna).
- " filiferum, n.sp. (Station 186, on Antedon variipinna).

OSTRACODA (Brady, Zool. pt. 3).

Macrocypris orientalis, Brady. (Station 187, 6 to 8 fathoms); obtained also at New Guinea. Recorded from Malay Archipelago.

Bairdia amygdaloides, Brady. (Station 187, 6 to 8 fathoms); for distribution see Station 162.

- " foveolata, Brady. (Station 187, 6 to 8 fathoms); for distribution see Station 33.
- ,, fortificata, n.sp. One specimen (Station 187, 6 to 8 fathoms); obtained at no other locality.

Cythere ovalis, n.sp. (Station 187, 6 to 8 fathoms); obtained at no other locality.

- " lubbockiana, n.sp. (Station 187, 6 to 8 fathoms); obtained at no other locality.
- " crispata, Brady. (Station 187, 6 to 8 fathoms); obtained also at Port Jackson and Hong Kong.
- " cancellata, Brady. (Station 187, 6 to 8 fathoms); obtained also at Station 172.