CAPE YORK.

Polyonyx obesulus (White), Miers. Four specimens (Station 186, 8 fathoms—three taken from interior of Hippospongia anomala); obtained at no other locality by the Challenger. Recorded from Madgica-Sima Islands and Australia.

Brachyura (Miers, Zool. pt. 49).

- Huenia proteus, De Haan. Four specimens (Station 186, 8 fathoms); obtained also in Arafura Sea. Recorded from China, Japan, and Philippines to Australia.
- Menæthius monoceros (Latreille), var. angusta, Dana. One specimen (Station 186, 8 fathoms); obtained also at Station 172.
- Chorilibinia gracilipes, Miers. One specimen (Torres Strait); obtained also at Station 188, 28 fathoms. Recorded from Australia and New Guinea.
- Schizophrys aspera (M.-Edwards). One specimen (Torres Strait); obtained at no other locality by the Challenger. Recorded from Indo-Pacific region.
- Pseudomicippa (?) varians, Miers. One specimen (Station 186,8 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.
- Lambrus affinis, M.-Edwards. One specimen (Torres Strait); obtained at no other locality by the Challenger. Recorded from Indo-Pacific region.
 - ", intermedius, Miers. One specimen (Station 187, 6 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and Corean seas.
 - (Aulacolambrus)hoplonotus, Adams and White. One specimen (Station 186, 8 fathoms); obtained at no other locality by the Challenger. Recorded from Indo-Pacific region.
 - ", () hoplonotus, var. granulosus, Miers. One specimen (Torres Strait, 3 to 11 fathoms); obtained at no other locality by the Challenger.
- Actæa granulata (Audouin). One specimen (Torres Strait); obtained at no other locality by the Challenger.
 - " hirsutissima (Rüppell). One specimen (Station 186, 8 fathoms); obtained at no other locality by the Challenger.
 - ,, hystrix, n.sp. One specimen (Station 186, 8 fathoms); obtained at no other locality.