

- Utriculus (Tornatina) acrobeles*, n.sp. (Station 186, 8 fathoms); obtained at no other locality. CAPE YORK.
- „ („), two species undetermined. (Station 186 and Torres Strait, 3 to 11 fathoms).
- Cylichna reticulata*, n.sp. (Stations 186 and 187, 8 and 6 fathoms); obtained at no other locality.
- „ *subreticulata*, n.sp. (Station 187, 6 fathoms); obtained at no other locality.
- „ (*Volvula*) *sulcata*, n.sp. (Station 186 and Torres Strait, 3 to 11 fathoms); obtained at no other locality.
- „ , three species undetermined. (Station 186 and Flinders Passage, 8 and 7 fathoms).
- Latirus* n.sp. (?). (Torres Strait, 3 to 11 fathoms).
- Assimineca* (?) sp. (Station 186, 8 fathoms).
- Leucorhynchia* (?) sp. (Station 186, 8 fathoms).

POLYPLACOPHORA (Haddon, Zool. pt. 43).

- Acanthopleura spiniger* (Sowerby). One specimen (Station 187, 6 fathoms); obtained at no other locality by the Challenger. Recorded from Australia and New Zealand to Philippines.
- Francisia spinosa* (Bruguière). (Station 187, 6 fathoms); obtained at no other locality by the Challenger. Recorded from Australia.

NUDIBRANCHIATA (Bergh, Zool. pt. 26).

- Chromodoris striatella*, Bergh. Three specimens (Station 186, 8 fathoms); obtained at no other locality by the Challenger. Recorded from Philippines and Torres Strait.
- Thordisa* (?) *clandestina*, n.sp. One specimen (Station 186, 8 fathoms); obtained at no other locality.

POLYZOA (Busk, Zool. pts. 30 and 50; Waters, pt. 79).

- Chlidonia cordieri* (Audouin). (Station 186, 8 to 11 fathoms); obtained at no other locality by the Challenger. Recorded from Atlantic and Mediterranean.
- Caberea lata*, Busk. (Station 186, 8 fathoms); obtained also at Station 190, 45 fathoms. Recorded from Australia and New Zealand.
- Carbasea cribriformis*, Busk. (Station 186, 8 fathoms); obtained also at Stations 188 and 190, 28 and 45 fathoms. Recorded from Australia and Cumberland Island. [Waters calls it *Flustra cribriformis* (Busk)].