

## AMBOINA.

- Madrepora effusa*, Dana. One specimen; obtained at no other locality by the Challenger.
- „ *gracilis*, Dana. Two specimens and fragments; obtained at no other locality by the Challenger.
- „ *appressa* (Ehrenberg). One specimen; obtained at no other locality by the Challenger.
- Turbinaria crater* (Pallas). One specimen; obtained also at Cape York and Wednesday Island.
- Montipora patula*, Verrill. One specimen; obtained at no other locality by the Challenger.
- „ *foliosa* (Pallas). Numerous specimens and fragments; obtained at no other locality by the Challenger.
- Porites palmata*, Dana. Obtained also at Philippines.
- Rhodaræa tenuidens*, n.sp. One specimen; obtained also at Philippines.
- Millepora intricata*, M.-Edwards and Haime. One specimen; obtained at no other locality by the Challenger.

## ASTEROIDEA (Sladen, Zool. pt. 51).

- Astropecten velitaris*, Martens. (15 to 25, or 100 fathoms?); obtained also at Admiralty Islands. Recorded from China.

## OPHIUROIDEA (Lyman, Zool. pt. 14).

- Ophiomusium simplex*, n.sp. (100 fathoms); obtained at no other locality.
- Ophionereis dubia* (Müller and Troschel). (100 fathoms); obtained also at Canary Islands (?) and Fiji (?).
- Ophiomyxa australis*, Lütken. (100 fathoms); for distribution see Station 109.

## ECHINOIDEA (Agassiz, Zool. pt. 9).

- Dorocidaris (Cidaris) bracteata*, n.sp. (15 to 25, and 100 fathoms); obtained at no other locality.
- Cælopleurus maillardi* (Michelin). (100 fathoms); obtained also at Stations 192 and 201.
- Salmacis dussumieri*, Agassiz. (100 fathoms); obtained also at Station 212, 10 to 20 fathoms.
- Clypeaster humilis* (Leske). (15 to 20 fathoms); obtained also at Station 212, 10 to 20 fathoms.
- Laganum putnami*, Barn. (15 to 25 fathoms); obtained at no other locality by the Challenger.
- Peronella decagonalis* (Lesson). (15 to 25 fathoms); for distribution see Station 172.