

ANOMURA (Henderson, Zool. pt. 69).

STATIONS 204
TO 204B.

Pagurus striatus, Latreille. One specimen (in shell of *Ranella fijiensis*, with attached spotted anemone); obtained also at Zebu reefs. Recorded from Mediterranean, North Atlantic, and Japan.

Eupagurus spinulentus, n.sp. One specimen (in shell of *Fusus niponicus*); obtained at no other locality.

Pagurodes piliferus, n.g., n.sp. One specimen; obtained also in Arafura Sea (?).

Paguroopsis typicus, n.g., n.sp. Two specimens; obtained at no other locality. Only species of the genus.

Galathea subsquamata, Stimpson. One specimen; obtained at no other locality by the Challenger. Recorded from Japan.

LAMELLIBRANCHIATA (Smith, Zool. pt. 35).

Pecten vitreus (Chemnitz). Obtained also at Stations 207, 209, 232, 307, 308, 310, and 311, 100 to 700 fathoms.

GASTEROPODA (Watson, Zool. pt. 42).

Fusus niponicus, Smith. Obtained at no other locality by the Challenger. Recorded from Nippon Island.

Pleurotoma (Surcula) deshayesii, Doûmet. Obtained at no other locality by the Challenger. Recorded from China.

Terebra (Myurella) mamillata, n.sp. Obtained at no other locality.

Ranella fijiensis, n.sp. Obtained also at Station 173.

Cassis (Bezoardica) wyvillei, n.sp. Obtained at no other locality.

Xenophora pallidula (Reeve). One specimen (beset with many other shells); obtained at no other locality by the Challenger. Recorded from Japan.

FISHES (Günther, Zool. pts. 6 and 57).

Champsodon vorax, Günther. Numerous specimens; obtained also at Stations 192 and 219.

Lioscorpius longiceps, n.g., n.sp. Obtained also at Station 192.

Lophius naresii, n.sp. Obtained also at Station 219, 150 fathoms, and Admiralty Islands.

Callionymus calauropomus, Rich. Obtained also at Station 162.

Monacanthus tessellatus, n.sp. Obtained at no other locality.

The Station-book records also an Actinian.